

The effects of elevated [CO₂] and decreased vapour pressure deficit on the nutrient status of maize and wheat plants under well-watered conditions

Kervin D. Prayag

PRYKER001

Thesis presented for the degree of Master of Science
in the Department of Biological Sciences

University of Cape Town

February 2020

Supervisors

Prof. Michael D. Cramer (University of Cape Town)

Prof. Brad S. Ripley (Rhodes University)



The copyright of this thesis vests in the author. No quotation from it or information derived from it is to be published without full acknowledgement of the source. The thesis is to be used for private study or non-commercial research purposes only.

Published by the University of Cape Town (UCT) in terms of the non-exclusive license granted to UCT by the author.

Plagiarism declaration

I know the meaning of plagiarism and declare that all the work in the document, save for that which is properly acknowledged, is my own. The thesis is submitted for the degree of Master of Science in the Department of Biological Sciences, University of Cape Town. It has not been for examination at any other university.

Signed: Signed by candidate

Date: 10/02/2020

Abstract

Increases in photosynthetic rates (A), biomass production and grain yield have been measured across a range of C_3 plants under elevated atmospheric $[CO_2]$ ('eCO₂'). However, decreases in the nutritional status of many C_3 plants growing at eCO₂ often occur concurrently with these increases. Several mechanisms have been proposed for these eCO₂-induced decreases, such as dilution effects due to enhanced carbohydrate production, down-regulation of photosynthesis, reduced root development, and decreased transpiration-driven mass flow delivery of nutrients. Reduced mass flow at eCO₂ is generally accepted as one cause for altered nutrient status in C_3 plants. However, eCO₂-induced reductions in mass flow remain understudied in C_4 plants, even though they account for about 18% of the total global net primary productivity and represent a large food source globally (e.g. maize and sorghum). This thesis investigated how mass flow reductions affect the nutrient status of wheat (C_3) and maize (C_4) plants. Reduced mass flow in both maize and wheat plants was induced with eCO₂ and by varying leaf-to-air vapour pressure deficits (VPD). I hypothesised that reduced mass flow at eCO₂ and at low VPD will negatively affect nutrient status in both the C_3 (wheat) and the C_4 (maize) species. In the first experiment, maize and wheat plants were grown at 400 and 800 ppm $[CO_2]$, in three well-watered soils, ranging from sandy to clayey, with and without fertilisation. In the second experiment, plants were grown at three VPD levels, namely 1.613 kPa, 0.773 kPa and 0.350 kPa, in well-watered soil and sand. In the latter experiment, to demonstrate the importance of mass flow, plants grown in sand were supplied nutrients in such a way that they had to rely exclusively on mass flow or diffusive processes (i.e. limited interception) for nutrient delivery to their rhizosphere. eCO₂ stimulated A on average by 22% in maize and by 50% in wheat, while stomatal conductance (g_s) and cumulative water loss (CWL) were respectively decreased by 35% and 31% in maize, and by 26% and 37% in wheat. eCO₂ reduced mass flow delivery of most nutrients on average by 32% in maize, and by 38% in wheat. The hypothesis that eCO₂-induced

reductions in mass flow negatively affect nutrient status in maize and wheat was however not supported. This was attributed to the well-watered conditions of the soils, which may have allowed for other processes (e.g. diffusion) to make up for the mass flow reductions. From 0.773 kPa to 0.350 kPa VPD, CW_L was decreased on average by 14% and 20% in the maize and wheat plants, respectively. A and g_s were little affected by VPD, but plants of both species always accumulated more biomass at 0.773 kPa. Consequently, there was little evidence to suggest that VPD-induced reductions in mass flow negatively affect nutrient status in maize and wheat. Reduced CW_L may have impeded root-to-shoot transport of ions and reduced dry biomass accumulation in the maize and wheat plants at 0.350 kPa (-40% and -22% on average respectively, relative to 0.773 kPa plants). Tissue [NPK] was also decreased (-13%, -41% and -47% respectively) in the 0.350 kPa VPD sand wheat plants, while increases in the proportion of finer roots may have alleviated effects of reduced CW_L on tissue [NPK] in the C_4 species. The findings from both experiments imply a decrease in the importance of mass flow for the delivery of nutrients to the rhizosphere under well-watered conditions. However, reductions in mass flow to a similar extent in both species at eCO_2 and low VPD — measured in the present study — suggest that under conditions of low water and nutrient availability, tissue nutrient concentrations could be negatively affected when transpiration is reduced.

Acknowledgements

I would like to first and foremost thank my main supervisor, Prof. Mike Cramer, who has provided impeccable supervision throughout my MSc. I will always strive to achieve the exemplary rigour and discipline that he applies to the scientific field. A massive thank you to Desmond Barnes, Calvin Hartnick and Dawood Hattas for their assistance in various technical aspects of my project. I cannot describe how much your contributions facilitated the execution of the growth experiments at UCT. Thank you to my co-supervisor, Prof. Brad Ripley, who has made me feel very welcome in the Botany Department at the Rhodes University since day one. Prof. Brad Ripley allowed me to use the Rhodes University Elevated CO₂ Facility, and without his collaboration, this project would not have been possible. Thank you to the technical officers (Dr. Riaan Strauss and Barry Hartley), and to the team of helpers at Rhodes University for their assistance, hard work and dedication throughout the eCO₂ experiment. Thank you to fellow students Simon Power, Kirsten Packer and Zander Venter for continuously providing valuable insights into my project. Dunja MacAlister, thank you for always being readily available to assist me with the root scans and with the IrGA. I am eternally grateful to Olivia Venter who always motivated me throughout my Master's, assisted with lab work, and continuously provided me with moral support. Thank you, Joshua Weiss, for your assistance during harvest. A massive thanks goes to Seth Musker for his extensive proofreading and wisdom throughout the Master's. Thank you to Ian Newton (Department of Archeometry, UCT) and to Raynette Topley (Elsenburg Labs, Stelleborsch) who promptly conducted various analyses on my soil and foliar samples. I also greatly appreciated the numerous tea/coffee breaks with my fellow postgrads and lecturers, namely Anthony Lowney, Gabriella Leighton, Hana Petersen, Mohammed Kajee, Dr. Robert Thomson and A. Prof. Colin Attwood.

This project was funded by the National Research Foundation.

Table of Contents

77	Plagiarism declaration.....	i
78	Abstract.....	ii
79	Acknowledgements	iv
80	Chapter 1	1
81	General Introduction	1
82	Hypotheses	11
83	Chapter 2	14
84	The effects of elevated atmospheric CO ₂ on the nutrient status of maize and wheat under	
85	well-watered conditions	14
86	Abstract.....	14
87	Introduction	15
88	Methods	20
89	Results	27
90	Discussion.....	29
91	Conclusion.....	33
92	List of figures.....	34
93	Figures	37
94	Chapter 3	47
95	The effects of low vapour pressure deficit on the nutrient status of maize and wheat under	
96	well-watered conditions	47
97	Abstract.....	47
98	Introduction	48
99	Methods	53
100	Results	59
101	Discussion.....	62

102	Conclusion	68
103	List of figures.....	68
104	Figures	71
105	Chapter 4	79
106	General discussion and synthesis	79
107	References	88
108	Appendices.....	113
109		
110		

Chapter 1

General Introduction

The global atmospheric CO₂ concentration ('[CO₂']') is increasing and is expected to reach 550 ppm by the middle of the century (IPCC, 2014). With this increase in atmospheric [CO₂], increases in photosynthetic rates, biomass production and grain yield are expected across a range of C₃ plants (Long et al., 2004; Ainsworth & Long, 2005; Ainsworth & Rogers, 2007; Leakey, 2009). These elevated [CO₂]-induced increases, however, often lead to altered nutritional status in many C₃ plants, as a result of overall decreases in concentrations of essential nutrients and minerals in the green tissue and grains of these plants. In a meta-analysis, Loladze (2014) reported between a 2% to 15% decrease (see Figure 1) in N, P, K, Ca, S, Mg, Fe, Zn and Cu and Mn in socio-economically important C₃ crops (such as rice, *Oryza sativa*, and wheat, *Triticum aestivum*). Similarly, a 10% decline in protein levels (see Figure 2) was observed in the edible parts of 18 rice cultivars grown at elevated [CO₂] (Zhu et al., 2018). In that same study, Zhu et al. (2018) also reported up to 30% decreases in B1, B2, B5 and B9 vitamin concentrations in the grains of these rice cultivars.




Figure 1: Changes in mean concentrations (%) of C, N, P, K, Ca, S, Mg, Fe, Zn, Cu, Mn, and in the ionome, in C₃ crop plants grown at an elevated [CO₂] concentration of 689 ppm, relative to C₃ plants grown at ambient [CO₂] (350 ppm). From Loladze (2014).




Figure 2: Changes (%) in protein concentration of 18 rice cultivars grown at elevated [CO₂] relative to ambient [CO₂]. From Zhu et al. (2018).

The decreases in essential minerals, proteins and vitamins in cereals that are socially and economically important across the globe can pose a potential threat to human nutrition (Myers et al., 2014). For example, the World Health Organization estimates that 25% of the global population suffer from Fe deficiencies, while 31% are Zn-deficient (Ezzati et al., 2004). Since crops such as wheat are also used in animal feed (in the form of grain or straw), CO₂-induced changes in nutrient status can also potentially affect the animal farming industry. For example, Porteaues et al. (2009) found that straw and grain quality in terms of crude protein was decreased in wheat plants grown at elevated [CO₂]. At an ecosystem level, the increased photosynthetic assimilation of CO₂ can lead to increased C:N ratios, with potential negative

impacts on nutrient cycling in natural ecosystems (Conroy, 1992). The physiological mechanisms resulting in these CO₂-induced changes in plant nutrient and mineral nutrition remain unclear but several hypotheses have been proposed: (1) altered root architecture and physiology (2) dilution of non-carbon compounds by increased carbohydrate production from enhanced photosynthesis (3) down-regulation of photosynthesis and decreased photorespiration, and (4) decreased stomatal conductance and transpiration rates resulting in reduced mass flow delivery of nutrients to the rhizosphere (Taub & Wang, 2008; Mcgrath & Lobell, 2013).

Several studies have reported changes in root architecture and physiology when plants are exposed to elevated [CO₂] (Berntson & Bazzaz, 1996; Pritchard & Rogers, 2000; BassiriRad et al., 2001). Rogers & Runion (1994) reported an increase in root length and number of root tips in crop plants correlating with high [CO₂], with the same observation being made in non-cultivated systems (Berntson & Woodward, 1992; Larigauderie et al. 1994). While an increase in the overall potential for nutrient and mineral uptake may be observed with longer and denser root systems, exploitation and uptake efficiency may be reduced (Fitter, 1987; Berntson, 1994). In addition, Pritchard & Rogers (2000) experimentally demonstrated that annual crops grown at elevated [CO₂] had more lateral roots versus primary roots, leading to more inefficient and shallower root systems, and ultimately reducing the volume of soil exploited (i.e. plants were not able to access resources in deeper soil layers). These changes in root architecture and physiology therefore hinder the efficiency of below-ground uptake of resources, which translates into reduced concentrations of a range of nutrients and minerals in plants at elevated [CO₂].

Changes in the nutrient status of plants grown at elevated [CO₂] have also been attributed to dilution, as a result of enhanced carbohydrate production due to increased photosynthetic assimilation of C (Eng et al., 1985; Wong, 1990; Gifford et al., 2000). This

process has been experimentally demonstrated in *Chrysanthemum morifolium* grown at elevated [CO₂]. After correcting for leaf starch, Kuehny et al., (1991) found no significant differences in the concentrations of Mn, Zn and Cu between *C. morifolium* plants grown at ambient and elevated [CO₂], and therefore attributed the uncorrected concentration decreases to dilution. The effect of dilution, however, may vary depending on the ability of plants to fix N₂. For example, Taub et al. (2008) reported a 1.5% decrease in N in soybean (a N₂-fixing species), while up to a 14% decrease was observed in non-N₂-fixing species. Furthermore, growing evidence suggests that dilution may not be the main process leading to decreased nutrient and mineral concentrations in plants exposed to elevated [CO₂] (Conroy, 1992; Taub & Wang, 2008; Feng et al., 2015). Poorter et al., (1997) experimentally demonstrated that mineral concentrations across a range C₃ crops, herbaceous and woody plant species, were consistently decreased at elevated [CO₂] even after correcting for total non-structural carbohydrates. Another argument advanced against the dilution hypothesis is that in plants grown at elevated [CO₂], all the elements should experience a degree of dilution, depending on their mobility and solubility (Zhao & McGrath, 2002; McGrath & Lobell, 2013). However, in a meta-analysis investigating the effects of elevated [CO₂] on the concentrations of macronutrients in the edible portions of food crops, Taub & Wang, (2008) reported that N experienced a greater decrease in concentration (-12%) compared to P (-5%) and K (-4%), suggesting that other physiological processes also play a role in the observed decreases of minerals and nutrients in plants exposed to high [CO₂].

Specifically, with regards to decreases of [N] reported in plants grown at high [CO₂], physiological processes affecting plant N acquisition, rather than dilution have been suggested. One such process is the inhibition of nitrate (NO₃⁻) photoassimilation, attributed to a decrease in photorespiration at elevated [CO₂] (Bloom et al., 2002; Searles & Bloom, 2003). NO₃⁻ reduction to NO₂⁻ occurs in the cytosol (Rufty et al., 1986) and requires NADH which is

generated from the OAA/malate and PGA/DHAP shuttles (Rathnam, 1978) that transports reductant from the chloroplast (Robinson & Giersch, 1987). At elevated $[\text{CO}_2]$, NADH reductant demand for CO_2 fixation can compete with NADH for the shuttle. Furthermore, the reduction of NO_2^- to NH_4^+ , the assimilation of NH_4^+ into amino acids, and the Calvin cycle all occur in the stroma and require ferredoxin. Elevated $[\text{CO}_2]$ also stimulates the Calvin cycle and can reduce the amount of ferredoxin available for NO_2^- reduction or NH_4^+ assimilation (Baysdorfer & Robinson, 1985; Peirson & Elliott, 1988). Two studies investigating N assimilation respectively in wheat and tomato plants reported a net decrease of NO_3^- assimilation at elevated $[\text{CO}_2]$ (Bloom et al., 2002; Searles & Bloom, 2003). In both studies, these decreases in NO_3^- assimilation were measured by calculating the assimilatory quotient (AQ), which is the ratio of net CO_2 consumption to net O_2 evolution: plants assimilating less NO_3^- show higher AQ values (Bloom et al., 1989). This decrease in NO_3^- photoassimilation can lead to a downregulation of shoot NO_3^- and the activities of the NO_2^- reductase enzyme, which in turn can potentially hinder the ability of crops to use NO_3^- as a source of nitrogen when exposed to conditions of high $[\text{CO}_2]$ (Bloom et al., 2002). Decreased NO_3^- assimilation can lead to an overall decrease in plant N concentration, and negatively affect yield and quality in wheat plants.

Another process affecting plant N acquisition at elevated $[\text{CO}_2]$ is the decrease in the activation, demand and production of the photosynthetic enzyme Rubisco (Long et al., 2004; McMurtrie et al., 2008; Taub & Wang, 2008), leading to a decrease in plant N demand at the leaf level. Rubisco is the most abundant leaf protein, accounting for up to 35% of total leaf N in C_3 plants (Evans & Seemann, 1989; Makino, 2003). A decrease in the production of this enzyme can therefore lower the overall N concentrations in plants. With this decrease in plant N investment in photosynthetic and photorespiratory structures comes an increase in photosynthetic plant N use efficiency (PNUE), as the N can be diverted into biomass

production (Taub & Wang, 2008). Although net losses of Rubisco have been shown in plants grown at elevated $[\text{CO}_2]$, this is not necessarily accompanied by decreases in net photosynthetic rates (Long et al., 2004), thus the increase in PNUE in plants growing in high $[\text{CO}_2]$. The decrease in the investment in photosynthetic enzyme Rubisco is due to the latter being carboxylated more readily and efficiently at high $[\text{CO}_2]$ levels (Davey et al., 1999; Stitt & Krapp, 1999; Gifford, Barrett & Lutze, 2000). When the Rubisco enzyme is saturated with CO_2 , binding of the former with the latter is increased, leading to a decrease in photorespiration. In addition, in these CO_2 -saturated conditions, the velocity of carboxylation by Rubisco is increased (Drake, González-Meler & Long, 1997). Webber, Nie & Long (1994) suggest that this situation can lead to the plants having the enzyme in excess in the photosynthetic apparatus and can represent an unnecessary investment of resources, particularly N, which could be diverted into the production of non-photosynthetic structures. However, a downregulation in the investment of photosynthetic enzymes can also decrease N availability to non-photosynthetic structures such as seeds and tubers as N is translocated to these structures from catabolised protein in the leaves (Fangmeier et al., 1999, 2002; Fangmeier, Chrost & Ho, 2000). Lower investment in the production of the Rubisco at high $[\text{CO}_2]$ can therefore lead to reductions in overall plant N content and concentration. Furthermore, changes in N concentrations in plants can lead to cascading effects in the concentrations of other minerals and nutrients in higher plants. For example, Cakmak, Pfeiffer & McClafferty (2010) demonstrated a positive correlation between N nutrition and the uptake of Zn and Fe in durum wheat. N affects root growth and the exudation of organic compounds by the root system (Marschner, 1995; Paterson et al., 2006). Thus, adequate N nutrition can firstly increase the direct interception of Zn and Fe by the root system, and secondly can increase the mobility of these two minerals in the soil as a result of increased exudation of organic compounds. In addition to downregulation of photosynthesis, changes in NO_3^- assimilation and changes in root

architecture, elevated [CO₂] can lead to an overall decrease in stomatal conductance and transpiration rate and can potentially represent an indirect effect of elevated [CO₂] on transpiration-driven mass flow delivery of nutrients to the rhizosphere.

Mass flow, along with diffusion and root interception are the processes that affect nutrient uptake via the root system. Uptake of nutrients by interception happens when roots are directly in contact with soil particles. This process is highly dependent on specific root length (root length/root mass). For example, high specific root length (SRL) leads to a high surface to volume ratio. This in turn maximises the amount of root surface in contact with the soil, and hence the potential of nutrient uptake (Larcher, 1995). Plants with high SRL have been found to show high uptake rates of N and P (Reich, Walters & Ellsworth, 1997; Comas, Bouma & Eissenstat, 2002). Diffusion acts along a concentration gradient resulting in dissolved nutrients in the soil solution moving from an area of high concentration to an area of low concentration until equilibrium is reached (Barber, 1995). Since there is continued uptake of nutrients by the root system, nutrients continuously diffuse towards the root surface. This process is particularly important in the acquisition of nutrients and minerals that are found in insoluble complexes, such as Fe (Mcgrath & Lobell, 2013). In contrast to diffusion, mass flow is the movement of water containing dissolved nutrients towards the roots as the plant transpires, allowing plants to access nutrients from beyond their rhizosphere (Barber, 1995). Transpiration-driven mass flow delivery of nutrients is considered as one of the most important processes affecting nutrient and mineral uptake below ground (Russell & Barber, 1960; Chapin, 1980; Lambers et al. 2008; Sellin et al., 2013). For example, Barber (1995) estimated that 79% N, 5% P, 18% K, 73% Ca, 88% Mg and 95% S were accumulated via transpiration-driven mass flow in maize plants growing in a fertile loam soil. It is well-known that reductions in stomatal conductance and transpiration rates at both the leaf and canopy scale occur when plants are exposed to elevated [CO₂] (Long et al., 2004; Bernacchi et al., 2006; Ainsworth & Rogers, 2007; Leakey

et al., 2009), and this can have effects on the transpiration-driven water fluxes for delivery and transport of minerals and nutrients in plants.

Decreased mass flow delivery of nutrients to the rhizosphere at elevated [CO₂] in C₃ plants have been reported in a few studies recently. In cottonwood (*Populus deltoides*) grown at elevated [CO₂], a 20% and 40% decrease in transpiration and stomatal conductance were respectively reported, accompanied by a 20% decrease in shoot [N], which was attributed to a decrease in transpiration-driven mass flow (McDonald, Erickson & Kruger, 2002). In a meta-analysis investigating the role of transpiration and altered nutrient allocation in the changes of nutrient status,, a significant correlation between decreases in concentrations of mobile nutrients — such as K, Ca and Mg — and a decrease in mass flow at high [CO₂] was found in wheat plants grown at eCO₂ (Mcgrath & Lobell, 2013). More recently, in a free-air CO₂ enrichment (FACE) experiment, the relationship between the below-ground uptake of N, K, S, Ca, Mg and Mn and transpiration rates in wheat plants at ambient and elevated CO₂ levels was demonstrated (Houshmandfar et al., 2018). It was found that the uptake of these elements was positively correlated with transpiration rates at both ambient [CO₂] and elevated [CO₂] (average R² = 0.86). A decrease in transpiration rate at elevated [CO₂] should logically lead to a decrease in the uptake of these resources, but the data in the latter study suggested that nutrient uptake per unit of water transpired was higher at elevated [CO₂], which might suggest that the plants were able to compensate for reduced mass flow. This uncertainty about the effects of elevated [CO₂] on water fluxes and nutrient uptake in plants shows the need for more research on this topic.

The idea that mass flow is reduced at elevated [CO₂] is generally accepted as one of the causes of altered nutrient status in C₃ plants such as wheat, rice and cottonwood (Taub & Wang, 2008; Leakey et al., 2009; Mcgrath & Lobell, 2013; Myers et al., 2014). This is because of the direct link between transpiration and CO₂ uptake in these plants. In C₄ plants however, the CO₂

concentrating mechanism leads to a relative decoupling between transpiration and CO₂ intake (Conroy, 1992; Bowes, 1993; Ghannoum et al., 2000; Taiz & Zeiger, 2010; Moore & Botha, 2014). As a result, there is some uncertainty about the relative effects that high [CO₂] will have on nutrient statuses in C₄ plants (Leakey et al., 2009; Myers et al., 2014). For example, at high [CO₂], little to no effect on NO₃⁻ photoassimilation due to changes in photorespiration, and on Rubisco activation and requirement, should be expected in C₄ plants. Although C₄ plants account for about 18% of the total global net primary productivity (Ehleringer et al., 1997) and represent a very large food source (e.g. maize and sorghum, *Sorghum bicolor*) globally (FAOSTAT), they remain understudied in the context of mass flow acquisition of nutrients (Loladze, 2014). Furthermore, Cramer, Hoffmann & Verboom (2008) experimentally showed that a C₃ plant, *Ehrharta calycina*, was able to regulate its transpirational flux in order to improve mass flow delivery of nutrients to its root system. In contrast, Rose et al. (2018) experimentally demonstrated that C₄ grasses were unable to regulate their transpirational fluxes under varying nutrient availabilities. The authors attributed this lack of regulation to less flexible WUEs in C₄ plants. Thus, under conditions of low nutrient availability, reductions in mass flow may more severely affect the nutrient status of C₄ plants.

In a study comparing the respective responses of cotton (C₃) and maize (C₄) plants at elevated [CO₂], a comparable decrease in transpiration rates was found between the C₃ and the C₄ species, while total N and protein content was only decreased in cotton (Wong, 1979). In addition, in a meta-analysis comparing the physiological responses of C₃ and C₄ plants at elevated [CO₂], stomatal conductance was decreased to a similar degree in C₃ and C₄ plants (Ainsworth & Long, 2005). In the same meta-analysis, it was found that photosynthetic carbon assimilation in both C₃ and C₄ plants was increased at elevated [CO₂], although that increase was greater in C₃ plants. The same conclusion was reached in a literature survey, wherein CO₂ assimilation was increased in both in C₃ and C₄ plants at elevated [CO₂], with the response

being stronger in C₃ plants (Cure & Acock, 1986). On the other hand, comparable increases in net CO₂ assimilation and total plant biomass were found between wild C₃ and C₄ grass species growing at eCO₂ (Wand et al., 1999). They argued that the photosynthetic mechanism in crop species such as maize and sorghum is already saturated at ambient [CO₂] levels, whereas in the wild grasses, saturation occurs at higher [CO₂], hence the comparable effects of [CO₂] on biomass and CO₂ assimilation in these grasses. This suggests that responses observed in cultivated systems may be very different in non-cultivated systems with regards to C₄ plants. This could have important implications for ecosystems management. Although significant decreases in transpiration rate and stomatal conductance at elevated [CO₂] have been reported in C₄ plants (Wong, 1979; Ainsworth & Long, 2005), other studies have reported little to no effect of elevated [CO₂] on the nutrient status of maize in well-watered conditions (Hocking & Meyer, 1991; Conroy, 1992; Manderscheid et al., 2014). More recently, in a FACE experiment, Erbs et al. (2015) found no significant differences in concentrations of N, P, K, Ca, Mg, S, Fe and Zn in both the aboveground biomass and in the kernels of well-watered maize plants grown at elevated [CO₂], even though stomatal conductance and transpiration rate were reduced. Studies investigating the physiological responses of C₄ plants to elevated [CO₂] (such as photosynthetic rate and growth stimulation) remain scarce (Sage, 1994; Ainsworth & Long, 2005; Loladze, 2014). Furthermore, the apparent contradictory results in terms of biomass accumulation in C₄ plants at elevated [CO₂] levels between studies highlights the need for further research on this topic.

Like elevated [CO₂], low vapour pressure deficit (VPD) also leads to decreases in transpiration rates in plants. Low VPD arises when relative air humidity levels are high. Since water evaporative demand is driven by the difference of water vapour pressure between leaf tissues and the atmosphere, in low VPD conditions, evaporative demand from the atmosphere is decreased, and as a consequence, transpiration rate of plants is decreased (Lihavainen et al.,

2016). Low VPD however promotes stomatal opening which allows for CO₂ to readily diffuse into the leaf mesophyll (Bunce, 1984; Gisléröd & Nelson, 1989), which subsequently leads to increased biomass accumulation. For example, when grown at low VPD, silver birch (*Betula pendula* Roth.) trees had higher biomass and higher levels of starch but accumulated lower concentrations of N and Fe (Lihavainen et al., 2016). Experiments where plants are grown at low VPD levels represent a good way to isolate the effects of decreased transpiration on nutrient concentrations and delivery in plants. Although elevated [CO₂] also leads to decreased transpiration rates in both C₃ and C₄ plants, it also has effects of other physiological process (i.e. photorespiration, Rubisco investment, root architecture and physiology), whereas low VPD levels seem to only affect transpiration rates. The fact that elevated [CO₂] and low VPD exert differential influences on transpiration in plants potentially allows for the contributors to CO₂ depression of the ionome to be better understood.

Hypotheses

In this thesis, I evaluated the effects of elevated [CO₂] and low VPD levels on water fluxes for resource delivery to the rhizosphere, biomass accumulation, and nutrient status in a C₃ crop (wheat) and a C₄ crop (maize). I hypothesised that elevated [CO₂] levels and low VPD levels will decrease tissue nutrient concentrations in both the maize and the wheat plants as a result of reduced transpiration-driven mass flow delivery of nutrients. To test this hypothesis, we divided the thesis into two data chapters:

In Chapter 2, I investigated the extent to which mass flow delivery is reduced in wheat and maize when grown at elevated [CO₂], as a result of decreased stomatal conductance and transpiration rate. Maize and wheat plants were grown at ambient or elevated [CO₂], in five well-watered soils of differing nutrient availabilities. Accumulation of nutrients by mass flow

was estimated by multiplying plant water use over the whole growth period by the respective concentrations of the nutrients in the soil solution. I then investigated the effects of elevated $[\text{CO}_2]$ on biomass accumulation and nutrient status in wheat and maize. I first hypothesised that elevated $[\text{CO}_2]$ will reduce tissue nutrient concentrations in wheat and maize, as a consequence of reduced mass flow delivery of nutrients to the rhizosphere. I then hypothesised that biomass accumulation will increase in wheat at elevated $[\text{CO}_2]$, but not in maize, leading to more pronounced decreases in nutrient concentrations in wheat plants as a result of dilution effects due to the different photosynthetic pathways of the two species. Finally, I hypothesised that mass flow-induced decreases in tissue nutrient concentrations in both species will be less pronounced with increasing soil nutrient availability.

In Chapter 3, I investigated the effects of varying VPD levels on transpiration-driven mass flow delivery of nutrients in maize and wheat. I expect the effects on nutrient status to be comparable between the two species. As in the elevated $[\text{CO}_2]$ experiment, accumulation of nutrients by mass flow was estimated by multiplying plant water use over the whole growth period by the respective concentrations of the nutrients in the soil solution. Maize and wheat plants were grown in two well-watered soil treatments, a clayey soil and a sand, at three VPD levels ranging from 1.613 kPa, 0.773 kPa and 0.350 kPa. In the sand treatment, to demonstrate the importance of mass flow, nutrients were introduced in such a way that the roots of the maize and wheat plants would not have direct access to the nutrients and would therefore have to rely solely on mass flow or diffusive processes (i.e. not interception) for nutrient delivery to the rhizosphere. I hypothesised that decreasing VPD levels will decrease tissue nutrient concentrations in the wheat and maize plants, as a result of decreased transpiration-driven mass flow delivery of nutrients to the roots. I then hypothesised that the negative effects of reduced transpiration-driven mass flow on the nutrient status of the plants will be more pronounced in

390 the sand treatment, where they had to rely exclusively on mass flow and diffusion for nutrient
391 delivery to their roots.

Chapter 2

The effects of elevated atmospheric CO₂ on the nutrient status of maize and wheat under well-watered conditions

Abstract

Increases in photosynthetic rates (A), biomass production and grain yield are expected across a range of C₃ plants with increases in atmospheric [CO₂] ('eCO₂'). However, these increases are often accompanied by decreases in the nutritional status of many C₃ plants. Several mechanisms have been proposed for these changes, such as dilution effects due to enhanced carbohydrate production, down-regulation of photosynthesis, reduced root development, and decreased transpiration-driven mass flow delivery of nutrients. Reduced mass flow at eCO₂ is generally accepted as one cause for altered nutrient status in C₃ plants while C₄ plants remain understudied in that context even though they account for about 18% of the total global net primary productivity and represent a large food source globally (e.g. maize and sorghum). This thesis investigated how mass flow reductions affect the nutrient status of wheat (C₃) and maize (C₄) plants. Reduced mass flow was induced with eCO₂ and it was hypothesised that reduced mass flow at eCO₂ negatively affect nutrient status in both the C₃ and the C₄ species. Maize and wheat plants were grown at 400 (ambient) and 800 ppm [CO₂] ('eCO₂'), in well-watered soils of differing nutrient availabilities. eCO₂ stimulated A on average by 22% in maize and by 50% in wheat, while stomatal conductance (g_s) and cumulative water loss (CW_L) were respectively decreased by 35 and 31% in maize, and by 26 and 37% in wheat. eCO₂ reduced mass flow delivery of most nutrients on average by 32% in maize, and by 38% in wheat. Nutrient status, however, remained unchanged in both species, attributed to the well-watered conditions of the soils, which may have allowed for other processes (e.g. diffusion) to make up for the mass flow reductions. The findings from this study suggest that negative effects of

eCO₂-induced reductions in mass flow on the nutrient status of plants may potentially be alleviated under well-watered conditions.

Introduction

Global atmospheric [CO₂] is on the rise and is expected to reach 550 ppm by the middle of the century even under the best mitigation scenarios (Ciais et al., 2013; Smith et al., 2016). With this increase in atmospheric [CO₂], increases in biomass accumulation and grain yield are expected in socio-economically important crops across the globe (Long et al., 2004; Ainsworth & Long, 2005; Ainsworth & Rogers, 2007; Leakey, 2009). These elevated [CO₂]-induced increases, however, are often accompanied by alterations in the nutritional status in many of these crops as a result of overall decreases in concentrations of nutrients essential for human and animal consumption in the green tissue and edible part of these crops, such as maize and wheat (Mcgrath & Lobell, 2013; Loladze, 2014).

Decreases in protein and mineral concentrations have been shown in wheat and maize plants grown under elevated [CO₂] ('eCO₂'), both in controlled indoor environments such as glasshouses and open-top chambers, as well as in experiments in open-field free-air CO₂ enrichment (Smith & Myers, 2019). The main mechanism proposed for these eCO₂-induced changes in plant tissue and grain stoichiometry is the dilution of non-carbon compounds due to increased production of carbohydrates (Loladze, 2014). For example, after correcting for leaf starch, no difference was found in the concentrations of Mn, Zn and Cu between *Chrysanthemum morifolium* grown at ambient and eCO₂ (Kuehny et al., 1991). Dilution effects at eCO₂ may vary depending on species. Jablonski, Wang & Curtis (2002) found no significant decrease in nutrient concentrations in a range of leguminous species at eCO₂ as a result of their ability to fix N₂, compared to a 15% decrease in non-leguminous species. However, this

mechanism alone cannot explain the totality of the eCO₂ effects as changes in nutrient status have been observed even when biomass accumulation was not stimulated (Feng et al., 2015).

Other physiological mechanisms have therefore been proposed, such as altered root architecture and physiology in both C₃ and C₄ plant species (BassiriRad et al., 1997; Pritchard & Rogers, 2000; BassiriRad, Gutschick & Lussenhop, 2001). An increase in root length and root tips has been shown in crop plants exposed to eCO₂ (Rogers & Runion, 1994). While these changes may increase overall nutrient uptake, soil exploitation and uptake efficiency may be reduced (Berntson, 1944; Fitter, 1987). Additionally, Pritchard & Rogers (2000) showed that in crops growing at eCO₂, the ratio of lateral to primary roots increased, leading to less efficient and shallower root systems, which may then impede on the plants ability to explore the deeper layers of the soil.

Down-regulation of photosynthesis and decreased photorespiration is another mechanism proposed for altered nutrient status in plants exposed to eCO₂ (Stitt & Krapp, 1999). Downregulation of photosynthesis at eCO₂ can lead to reductions in the activity, demand for, and content of the photosynthetic enzyme Rubisco, which subsequently will decrease plant leaf N demand (Long et al., 2004; McMurtrie et al., 2008; Taub & Wang, 2008). Since Rubisco is the most abundant leaf protein and makes up to 35% of total leaf N in C₃ plants, a decrease in the production of this enzyme at eCO₂ can lead to a decrease in the total leaf N content (Evans & Seemann, 1989; Makino, 2003). In C₄ plants, such as *Sorghum bicolor*, it was found that ratios of Rubisco to PEP-Carboxylase remained constant at eCO₂ during leaf development (Cousins et al., 2003). However, in the same study, it was shown that the total activity of Rubisco was decreased in older leaves, suggesting that C₄ plants may also down-regulate photosynthetic activity at eCO₂, which can lead to reductions in leaf N demand and leaf N content.

Decreased stomatal conductance and transpiration rates at eCO₂ can also lead to reduced nutritional value of crops (Taub, Miller & Allen, 2008; Mcgrath & Lobell, 2013). These decreases can lead to reductions in overall plant water use, and subsequently in decreased transpiration-driven mass flow delivery of nutrients to the root systems of plants. Mass flow, together with diffusion and root interception are important processes affecting below-ground nutrient uptake in plants (Barber, 1995; Cramer, Hoffmann & Verboom, 2008). For example, (Barber, 1995) estimated that mass flow contributed to 79% N, 5% P, 18% K, 73% Ca, 88% Mg and 95% S in maize plants growing in a fertile loam. Cramer, Hoffmann & Verboom (2008) experimentally demonstrated that *Ehrarta calycina*, a C₃ grass, has the ability to regulate its transpirational flux in order to increase mass flow delivery of nutrients to its roots, when soil nutrient availability is low. However, this regulation was not found in C₄ grasses, wherein no increase in transpiration was measured at low soil nutrient availabilities (Rose et al., 2018). The authors attributed this lack of response to less flexible water-use efficiencies (WUE) in C₄ species. Since decreases in stomatal conductance and transpiration rate are measured in both C₃ and C₄ plants species at eCO₂ (Ainsworth & Long, 2005), the apparent inability of C₄ species to regulate transpirational fluxes at differing below-ground nutrient availabilities may put them at a greater disadvantage than C₃ plants in future climate change scenarios. Decreased mass flow of nutrients at eCO₂ is generally accepted as one of the causes of decreased nutritional value in C₃ crops such as wheat and rice (Taub & Wang, 2008; Mcgrath & Lobell, 2013; Myers et al., 2014). Due to the fact that their photosynthetic mechanism is already saturated at current CO₂ levels (Ziska & Bunce, 1997; Ainsworth & Rogers, 2007), some uncertainties remain about how eCO₂-induced decreases in mass flow will affect nutrient concentrations and content in C₄ crops such as maize (Leakey et al., 2009; Myers et al., 2014).

Often, availability of nutrients and water interact with and dictate the effects of eCO₂ on plant biomass and nutrient status (Körner, 2006). For example, it has been argued that

negative effects of eCO₂ on grain protein concentration in wheat would be alleviated with increased availability of N (Evans, 1993; Pleijel et al., 1999). However, in a more recent study, it was shown that grain protein concentration in this C₃ crop significantly decreased at eCO₂ regardless of N fertilisation (Pleijel et al., 2019). Additionally, it also been shown that downregulation of photosynthesis in C₃ plants happens to a greater extent in low N environments compared to high N environments (Stitt & Krapp, 1999; Ainsworth & Rogers, 2007). In maize grown at eCO₂, it was found that yield and leaf physiological capacity remained unchanged across a range of soil N fertility, provided the plants were not water-stressed (Markelz, Strellner & Leakey, 2011). In the same study, decreases in leaf number, leaf area index, biomass, and yield were recorded at low N availability in the maize plants grown at both ambient and eCO₂, suggesting that nutrient stoichiometry was not affected by variation in atmospheric [CO₂].

With regards to interactions between soil water availability and eCO₂, it was found that stomatal conductance decreased by 51% in droughted wheat plants grown at eCO₂, compared to a 41% decrease under well-watered conditions (Li, Kang & Zhang, 2004). Similarly, in maize plants, stomatal conductance at eCO₂ was reduced by 57% in droughted conditions, and by 33% in well-watered conditions (Markelz, Strellner & Leakey, 2011). Since mass flow is directly related to plant transpiration, differing watering regimes are likely to influence eCO₂-induced decreases in mass flow delivery of nutrients, and subsequently influence eCO₂ effects on nutrient status. Increased water supply may thus alleviate the negative effects of eCO₂-induced decreases in mass flow on nutrient status, especially in C₄ plants with less flexible WUEs than C₃ plants (Ainsworth & Rogers, 2007). While there are some similarities as to how eCO₂ affects stomatal conductance and transpiration rates in C₃ and C₄ plants, the eCO₂ effects on photosynthesis differ (Ainsworth & Rogers, 2007). Stimulation of photosynthesis in C₃ crops grown at eCO₂ seem to occur under a range of watering regimes (Kimball, 2016). In

maize plants however, stimulation of photosynthesis at eCO₂, and subsequent increase in biomass, was generally found only under dry conditions, as a result of improved water-use efficiency (Leakey et al., 2009; Markelz, Strellner & Leakey, 2011; Erbs et al., 2014). Erbs et al. (2014) argue that eCO₂ could mitigate the impacts of drought stress on nutritional quality of maize plants. However, increases in biomass accumulation, combined with reductions in stomatal conductance (and consequently reductions in mass flow delivery of nutrients), could severely decrease the nutritional status of these C₄ plants under conditions of drought and eCO₂.

Since decreases in stomatal conductance and transpiration rates have been reported in both maize and wheat at eCO₂ (Ainsworth & Rogers, 2007; Leakey et al., 2009), I hypothesised that these reductions will lead to a decrease in tissue nutrient concentrations in both the C₃ and the C₄ crop as a result of reduced mass flow delivery of nutrients. I then hypothesised that reduced root development at eCO₂ in both species will also contribute to decreases in tissue nutrient concentrations. Since photosynthesis is mostly likely to be stimulated by eCO₂ in plant C₃ species compared to C₄ plant species under well-watered conditions (Ainsworth & Rogers, 2007), I hypothesised that nutrient concentrations will be decreased to a greater extent in wheat compared to maize, as a result of dilution effects due to enhanced carbohydrate production. Finally, I hypothesised that the negative effects of eCO₂-induced reductions in mass flow on the nutrient status of both crop species will be lessened with increasing soil nutrient availability (Körner, 2006; Mcgrath & Lobell, 2013). To test these hypotheses, maize and wheat plants were grown at two CO₂ levels, under well-watered conditions, and in soils with differing nutrient availabilities, and changes in nutrient concentrations and contents were measured for several important nutrients.

Methods

Plant germination and growth conditions

Wheat (*Triticum aestivum* c. SST015, Kaap Agri, Malmesbury, 7300, South Africa) and maize (*Zea mays* c. CAP 122-60, Capstone Seeds, Howick, 3290, South Africa) were grown from seeds soaked in aerated water for 24 h and 48 h, respectively, before being sown in trays of vermiculite. The trays were kept moist until the emerging seedlings were established. The seedlings were transplanted into pots, each containing 7 kgs of soil. Three soils of varying clay content were used in this experiment: a top soil with high clay content (hereafter referred to as 'Top Soil'), sand with low clay content (hereafter referred to as 'Sand'), and a mix of sand (70% by mass) and top soil (30%), with moderate clay content (hereafter referred to as 'MIX').

Plants growing in Top Soil and MIX were subjected to two fertilisation treatments: an unfertilised treatment and a treatment fertilised with 25 g of Protek (Farmwox (Pty) Ltd Reg. No.: B4652) pellets containing (mmol.g⁻¹) 3.43 N; 0.32 P; 0.92 K; 0.0003 Cu; 0.016 Fe; 0.004 Mn and 0.00009 Zn. The fertilised treatments of Top Soil and MIX are hereafter respectively referred to as 'Top Soil NPK' and 'MIX NPK'. Plants growing in Sand were always subject to fertilisation, with 25 g of Protek pellets and a modified Long-Ashton solution containing only 1.5 mM MgSO₄.7H₂O, 4 mM CaCl₂.2H₂O, 0.1388 mM H₃BO₃, 0.0208 mM MnSO₄.4H₂O, 0.0023 mM ZnSO₄.7H₂O, 0.0033 mM CuSO₄.5H₂O, 0.0002 mM Na₂MoO₄.2HO and 0.0899 Fe EDTA (Hewitt, 1966). The Sand pots were first supplied with excess water before being supplied with 200 mL of solution twice weekly. The soil fertilisation treatments (hereafter referred to as 'soil treatments') were chosen so as to have a nutrient availability gradient. Thus 'Top Soil', 'Top Soil NPK', 'MIX', 'MIX NPK' and 'Sand' were treated as distinct soil treatments with different nutrient availabilities (see Table 1 below).

These soil treatments were replicated so as to have wheat and maize plants subject to them at either 400 ppm (hereafter referred to as ‘ambient’), or at 800 ppm (hereafter referred to as ‘eCO₂’). The plants growing at ambient CO₂ were randomly distributed between two open-top chambers into which the atmospheric [CO₂] was set at 400 ppm. Similarly, the plants growing at eCO₂ were randomly distributed between two open-top chambers into which the atmospheric [CO₂] was set at 800 ppm. The experiment was thus conducted using a split-plot experimental design, with the four open-top chambers being the whole-plots, and the split-plots being the different soil treatments in which the plants were growing. The experiment was conducted at the Rhodes University Elevated CO₂ Facility (Rhodes University, Grahamstown, South Africa, 33°18'41.2"S 26°30'33.4"E).

Field capacity and water use

Prior to running the experiment, the respective field capacities of the three soil types were determined. This was done by supplying excess water to pots containing 200 g of soil that had been oven-dried 7 d at 70°C to remove any moisture. The pots were covered with plastic sheets to prevent evaporation and were allowed to settle for 24 h. Field capacity of the different soil types was then determined as the dry weight percentage of the difference between the dry weight of the soil and the wet weight 48 h after they had settled (Kirkham, 2005). The field capacities were 28%, 17% and 14 % for Top Soil, MIX and Sand, respectively. Throughout the experiment, the soils were kept between 70-100 % of their respective field capacities by gravimetric measurement of water content. These field capacities were chosen to allow unrestricted mass flow (Marschner & Rengel, 2011). For each soil type, 10 additional control pots filled with the same amount of soil (7 kg) were randomly distributed in the two open-top chambers set at ambient, and in the two open-top chambers set at eCO₂. These control pots were subject to the same watering regime as the bags with the plants. This was done so as to gravimetrically estimate the cumulative water use per plant throughout the growth experiment.

Soil nutrient availability (shown in Table 1)

A soil column was constructed from a 20-mL syringe with the plunger removed, and a disc of Whatman No. 1 filter paper was placed at the bottom. The filter paper was topped with glass wool to prevent loss of sediment. Oven-dried soil was filled to the 20 mL mark and water added to field capacity. The soil column was inserted through a hole in the lid of a 50-mL Falcon centrifuge tube, placed in the tube, and centrifuged (Eppendorf Centrifuge 5810R) at 150 g for 2 min. The volume of water collected in the centrifuge tube was taken as a relative measure of the mass flow displaceable soil solution. The mass flow technique using low-speed centrifugation was a slight modification of the gravity method described by (Barber, 1995). The elemental determination in the water extracts were performed by the Elsenburg Laboratory (Western Cape Department of Agriculture, Stellenbosch, South Africa, 33.845259 S; 18.834722 E). The concentrations of P, K, Ca, Cu, Fe, Mn and Zn were determined directly using a Radial ICP Spectrometer ICAP 7600 (Thermo Fisher Scientific, Sunnyvale, CA, USA), while N was determined on a BR-71394 DA Gallery Analyzer (Thermo Fisher Scientific, Sunnyvale, CA, USA). Concentration of most nutrients in water extracts from Top Soil NPK, MIX NPK and Sand were generally higher compared to water extracts from Top Soil and MIX (Table 1).

Table 1: Concentrations (mg l⁻¹) of N, P, K, Ca, Cu, Fe, Mn and Zn in soil solution extracted from the five soil types in which the wheat and maize plants were grown. Different lower case letters indicate the significant differences amongst soil treatments as determined by Tukey's post-hoc comparisons.

	Top Soil	MIX	Top Soil NPK	MIX NPK	Sand
Total inorganic N	10.68 (a)	9.29 (a)	32.63 (b)	46.57 (c)	49.04 (c)
Phosphorus	1.00 (a)	1.00 (a)	4.20 (b)	6.80 (c)	14.00 (d)
Potassium	17.80 (a)	14.80 (a)	29.00 (b)	37.00 (c)	42.40 (d)
Calcium	78.00 (a)	68.20 (a)	107.00 (b)	117.80 (b)	128.80 (b)
Copper	0.05 (a)	0.04 (a)	0.05 (a)	0.06 (a)	0.07 (a)
Iron	2.60 (d)	1.46 (bc)	1.68 (c)	0.88 (ab)	0.65 (a)
Manganese	0.04 (a)	0.05 (a)	0.06 (b)	0.09 (c)	0.09 (c)
Zinc	0.87 (b)	0.61 (b)	0.87 (b)	1.58 (c)	0.16 (a)

Gas exchange measurements

All plant gas exchange measurements were conducted on fully expanded leaves using a LI-6400 (Li-Cor, Lincoln, NE, USA) Portable Photosynthesis System equipped with 2 x 3 mm broadleaf cuvette and an integrated red/blue LI-6400-02B LED light source (Li-Cor, Lincoln, NE, USA). I set the [CO₂] in the cuvette to 400 or 800 ppm depending on the CO₂ level at which the plants being measured were exposed. Relative humidity was set to between 55-65% and block temperature was set to match the temperature of the day of which the measurements were being taken. The variables measured were photosynthetic rate, stomatal conductance and transpiration rate.

Plant harvest

The wheat and maize plants were harvested after 7 weeks of growth. The soil was carefully washed away with water on a 2-mm sieve to minimise root loss. The plants were separated into root, stem and leaves, the surface dried with paper towel, and immediately weighed to obtain fresh weight (stem and leaves were later combined to obtain above-ground biomass). The leaves were then scanned with a flatbed scanner (Canon CanoScan LiDE 220) so as to obtain the leaf area and thus the specific leaf area (SLA) of each plant. The harvested plants were then oven-dried at 70°C for 48 h after which dry weight was determined. A representative subsample of the root system of each plant (Vandamme et al., 2013) was used to determine average root diameter (mm) and total root surface area (m²) using a STD4800 scanner and WinRHIZO version 2013a (Regent Instruments, Canada). Specific root length (SRL, m g⁻¹) was calculated as the total root length (m) divided by the root dry weight (g). The root characteristics were converted to whole-root results by multiplying by their respective conversion factors to 100% of the root.

Mass flow delivery and contribution

Mass flow delivery of each nutrient was calculated as the cumulative amount of water transpired by a plant throughout the growth experiment multiplied by the concentration of that nutrient in the soil solution. Since delivery of a nutrient does not necessarily equate to acquisition, mass flow contribution to tissue concentration of that nutrient was then calculated as a percentage of the total concentration of that nutrient in the plant. The rest was assumed to have been acquired by diffusion and/or root interception (Barber, 1995).

639 *Foliar elemental analyses*

640 The dried foliar and root samples were ground to a fine powder using a grinder or a ball-mill
641 (MM200, Retsch, Germany). The powdered samples were placed in Perspex rings sealed with
642 4 µm Polypropylene thin film (Chemplex Industries Inc, Florida, USA) and introduced to a
643 SPECTRO XEPOS XRF spectrometer (SPECTRO, AMATEK materials analysis division,
644 Kleve, Germany). Analyses were conducted using the X-LabPro 5 software, which
645 incorporates the universal ‘Turbo Quant Powders’ method. The instrument was calibrated by
646 using a certified standard GBW07312 (National Research Center for CRMs, Beijing, China),
647 for which elemental concentrations were obtained from NOAA Technical memorandum NOS
648 ORCA 68 (1992). Only the elements that were within the machine’s detection limits were
649 included.

650 Total N concentration of the powdered foliar and root samples were determined using
651 mass spectrometry. Samples were weighed into tin capsules (5 x 9 mm; Sántis Analytical,
652 Teufen, Switzerland, with *ca.* 2 mg of foliar powder used for analysis). The samples were
653 combusted in a Flash 2000 organic elemental analyser and the gases passed into a Delta V Plus
654 isotope ratio mass spectrometer (IRMS) via a Conflo IV gas control unit (all from Thermo
655 Scientific, Bremen, Germany). In-house standards and one IAEA (International Atomic Energy
656 Agency) standard (USGS25) were used to calibrate the results. Nitrogen concentration was
657 expressed relative to atmospheric nitrogen (Evans, 2001). Ratios of the different elements were
658 obtained by dividing their respective concentrations by their carbon concentrations.

659 *Data analyses*

660 All statistical analyses were performed using R Statistical Software (R Core Team, 2019).
661 Comparisons of all the response variables were made using linear models with CO₂ level and
662 soil treatment as explanatory variables, fitted as interactions, or as separate main effects if the

interaction was not significant. Random effects were included in the fitted models to account for the non-independent structure of the experimental design, since plants growing in one open-top chamber are more likely to experience similar growing conditions compared to those growing in another open-top chamber – that is even if they were growing in the same soil treatment and were exposed to the same CO₂ concentration in two different open-top chambers. However, the random effects were found to be non-significant in all instances, potentially demonstrating the homogeneity in terms of the controlled environment that the open-top chambers at the Rhodes University Elevated CO₂ Facility can provide. Consequently, the random effects were dropped from the models, and the data was analysed in bulk (in terms of chambers). The concentrations of N, P, K, Ca, Cu, Mn, Fe and Zn in the water extracts from the different soil treatments were also compared using linear models, with concentrations of the elements as the response variables, and soil treatment fitted as the explanatory variable. Analyses of variance were done on all the fitted linear models to determine the significance of the main effects using the ‘car’ package (Fox & Weisberg, 2011). Where main effects significantly explained variation, pairwise comparisons (Tukey post-hoc tests) of group level means were then performed using the package ‘emmeans’ (Lenth, 2018) to determine which soil treatments showed a significant effect of [CO₂]. To test the relationship between mass flow delivery and nutrient concentration, the data were pooled in terms of soil treatments to fit linear models with tissue nutrient content as the response variable and mass flow delivery of nutrients as the explanatory variable. The same was done to generate correlation plots between these two variables. Boxplots and correlation plots were all generated using the ‘ggplot2’ package (Wickham, 2016).

Figures 1 to 10 and Appendices 1 to 12 show the effects of soil treatment and [CO₂] on morphological, physiological and nutrient variables in the maize and wheat plants. The ANOVA results indicating the significance of each factor are shown in each plot, with

interaction terms excluded if they were not significant ($p > 0.05$). Note that a significant interaction term implies that the effect of $[\text{CO}_2]$ varied between soil treatments.

Results

Total biomass, root biomass, shoot-to-root ratio and root characteristics

eCO_2 did not have any significant effect on total biomass, root biomass or shoot-to-root ratio in maize plants (Fig. 1). eCO_2 increased total biomass and root biomass in wheat plants, but this was dependent on soil treatments. Total biomass was significantly higher in wheat plants at eCO_2 in MIX NPK, and there was no difference in tissue nutrient concentrations in both species exposed to the two CO_2 levels in the other soil treatments. Root biomass was higher at eCO_2 in MIX NPK and Top Soil NPK, with no difference between the two CO_2 levels in the other soil treatments. There was no difference in shoot-to-root ratio in wheat between the two CO_2 levels in any of the soil treatments.

There was no effect of eCO_2 on average root diameter, total surface area and specific root length in maize plants within each soil treatment (Fig. 2). Wheat plants grown at eCO_2 showed increased average root diameter in all soil treatments, but no differences in total surface area. Wheat plants grown at eCO_2 generally showed decreased specific root length, but this effect was dependent on soil treatment, with the only significant reduction occurring in plants grown in MIX NPK.

Gas exchange and water use

Stomatal conductance and transpiration rate were decreased in both maize and wheat plants grown at eCO_2 in all soil treatments (Fig. 3). eCO_2 increased photosynthetic rate in wheat plants in all soil treatments. There was an overall significant effect of $[\text{CO}_2]$ on photosynthetic rate in

maize plants, but post-hoc results showed no difference between the two [CO₂] levels within each soil type. Cumulative water use (i.e. total water loss) per plant was decreased in maize plants grown at eCO₂ in all soil treatments (Fig. 4). This was not the case in wheat where cumulative water use per plant was decreased in Top Soil, MIX and Sand, but not in Top Soil NPK and MIX NPK.

Mass flow delivery and contribution

The eCO₂-induced reduction in cumulative water use resulted in decreased mass flow delivery of N, P, K, Ca, Cu, Mn, Fe and Zn in maize plants in all soil treatments (Fig. 5, Appendices 1 and 2). Similarly, in wheat, eCO₂ plants had reduced delivery of all the nutrients by mass flow, but this was dependent on soil treatment, since in Top Soil NPK and MIX NPK, mass flow delivery was not reduced. Maize plants grown at eCO₂ also had decreased mass flow contribution to tissue content of most nutrients (except Fe), but this was dependent on soil treatment with the only significant reduction occurring in plants grown in Sand (Fig. 6, Appendices 3 and 4). In wheat, eCO₂ plants showed significantly reduced mass flow contribution of most nutrients in all soil treatments.

Nutrient concentrations, contents, ratios and, tissue nutrient content versus mass flow delivery

Despite marked changes in variables relating to nutrient uptake, there were no differences in tissue concentrations of most of the nutrients in wheat and maize plants grown at the two CO₂ levels (Fig. 7, Appendices 5 and 6). Tissue nutrient contents were also little affected by eCO₂ in maize (Fig. 8, Appendices 7 and 8). In wheat, contents of all the elements, except for Fe, were increased at eCO₂ in the MIX NPK soil. Additionally, contents of N, Ca, Mn and Fe were increased at eCO₂ in the Top Soil NPK treatment.

Linear regressions revealed significant positive correlations between N, P and K tissue content and amount of these nutrients delivered by mass flow, in both maize and wheat (Fig. 9). Additionally, the intercepts (i.e. when 0 mg N, P, and K delivered by mass flow) were higher at eCO₂ in both maize and wheat for N, P and K. The correlation between amount of Fe delivered by mass flow and tissue Fe content revealed a significant negative correlation in wheat plants at both ambient and eCO₂ (Appendix 10). For the other nutrients, depending on the species, correlations were either not significant at ambient, eCO₂ or both [CO₂] levels (Appendices 9 and 10).

Wheat plants grown at eCO₂ showed increased C:N and C:P ratios in all soil treatments except MIX (Fig. 10). Increased C:K, C:Ca, C:Mn and C:Zn ratios were observed in wheat plants grown in the MIX NPK soil (Fig. 10, Appendices 11 and 12). Changes in these ratios can indicate dilution effects of some elements as a result of enhanced carbohydrate production at eCO₂. In contrast to wheat, maize plants grown at eCO₂ did not have different ratios.

Discussion

In agreement with the literature, eCO₂ reduced stomatal conductance and transpiration rates in both maize and wheat plants (Long et al., 2004; Bernacchi et al., 2006; Ainsworth & Rogers, 2007; Leakey, 2009). Cumulative water use per plant was also decreased in both species at eCO₂, except for wheat plants growing in Top Soil NPK and MIX NPK. As such, transpiration-driven mass flow delivery of nutrients was significantly decreased at eCO₂, except in the wheat plants growing in Top Soil NPK and MIX NPK. However, there was little to no decrease in tissue nutrient concentrations at eCO₂ in either of the crops, regardless of soil treatment. Since no decrease in mass flow delivery of nutrients was recorded in the wheat plants growing in Top Soil NPK and MIX NPK, adequate supply of nutrients can explain the lack of change in their

tissue nutrient concentration at eCO₂. The following question therefore arises: why then were there no decreases in tissue nutrient concentrations in both maize and wheat at eCO₂ even in the instances where mass flow delivery of nutrients was significantly decreased?

The amount of a nutrient displaced by mass flow is strongly dependent on its solubility in soil water (Barber, 1995). For example, Ca is highly soluble and is often delivered in excess to the rhizosphere (Lambers, Chapin & Pons, 1998; Zhao & McGrath, 2002; Cramer & Hawkins, 2009). Fe complexes are generally less soluble, making the diffusive process more important than mass flow for the transport of these nutrients (Mcgrath & Lobell, 2013). Additionally, it is important to make the distinction between nutrient delivery and acquisition, whereby delivery by mass flow does not necessarily equal acquisition. In agreement with the literature (Cramer & Hawkins, 2009), it was found that Ca was delivered in excess of physiological requirements at both ambient and eCO₂ in both maize and wheat plants. As such, the decreases in mass flow contribution of Ca measured at eCO₂ in both the C₃ and C₄ crop species in the present study did not negatively impact their tissue Ca concentration or content. Furthermore, negative relationships (and non-significant in maize) were found between tissue Fe content and mass flow delivered Fe in both species, suggesting that diffusive processes or root interception may have been more important for the delivery of this nutrient to the rhizosphere. Thus, even though a decrease in mass flow contribution of Fe was measured in some soil treatments in the wheat plants at eCO₂ (no decrease in maize), no decrease was measured in the concentration of this nutrient in either of the species.

Mass flow and diffusive processes are strongly dependent on water flow in the soil (Barber & Cushman, 1981; Brouder & Volenec, 2008), and so is their relative importance (Cramer, Hoffmann & Verboom, 2008). Thus, the well-watered conditions (between 70-100% field capacity) to which the maize and wheat plants were subject, at both ambient and eCO₂, may have allowed for these two processes to adequately and interchangeably transport nutrients

to the root systems, regardless of soil nutrient availability, even though cumulative water use per plant and mass flow contribution of most of the elements were reduced at eCO₂. Schmidt et al (2004) showed that although an increase in the soil solution concentrations of several elements can occur during and after a drought event, the flux of these elements is generally lower due to decreased water flow in water-stressed environments. The well-watered status of all the soil treatments in the present study can therefore explain the lack of negative effects of eCO₂-induced reductions in mass flow delivery of nutrients on the nutrient status of the plants. Additionally, eCO₂ tends to significantly affect mass flow while having small effects on diffusive processes (Mcgrath & Lobell, 2013). Therefore, reduced mass flow induced by eCO₂, if combined with conditions of water stress, could lead to a decrease in the tissue nutrient concentrations of these two crops. When looking at the relationship between mass flow delivery and tissue contents of N, P and K, the significantly higher contents of N, P and K at eCO₂ in the absence of mass flow (at intercept) in both species, suggests a decrease in the importance of this process, in favour of other processes such as diffusion or interception for the delivery and acquisition of these nutrients. These results may explain the patterns reported in the study by Kimball (2006), in which decreases in tissue nutrient concentrations in wheat and cotton grown at eCO₂ were alleviated in well-watered treatments (4% decrease in foliar N concentration as opposed to 9% under limited water supply). Thus, the hypothesis that eCO₂-induced decreases in mass flow delivery of nutrients negatively affect the nutritional value of maize and wheat is not verified in this study, possibly as a result of the well-watered conditions of all the soil treatments.

When considering the altered root architecture hypothesis, the lack of changes in specific root length and total surface area in both maize and wheat at eCO₂, in addition to constant flow of water below-ground, may have allowed for adequate nutrient delivery and uptake, and can further explain the lack of difference in tissue nutrient concentrations between

the two CO₂ levels in both species. With regards to root diameter, BassiriRad et al. (1997) demonstrated that C₄ plants were able to increase the proportion of finer roots and maintain adequate N and P concentrations at eCO₂, while the concentrations of these nutrients were decreased in C₃ plants. Additionally, Gillespie (1989) has demonstrated that finer roots can more efficiently extract nutrients such as N, P and K from the soil. In the present study, root diameter was increased in the wheat plants grown at eCO₂, thereby potentially leading to less efficient root systems for belowground nutrient uptake in the C₃ crop (Luo et al., 2013). Nonetheless, nutrient concentrations remained unchanged in both the C₃ and the C₄ species. In a recent study done by Houshmandfar et al. (2018), it was shown that eCO₂ can increase the nutrient uptake per unit of water transpired in wheat, attributed to increased root growth in the earlier stages of growth. Data on root growth at different stages of growth were not available in the present study, but it is possible that changes in root ontogeny allowed the eCO₂ wheat plants to maintain a similar nutrient uptake rate to that of their ambient CO₂ counterparts. Therefore, the hypothesis that altered root architecture at eCO₂ will lead to decreases in nutritional value of the C₃ and C₄ crop is also not verified in this study.

eCO₂ significantly increased photosynthetic rate in wheat within each soil treatment, but this was not observed in maize, which is expected due to the differing C₃ and C₄ photosynthetic pathways. The photosynthetic mechanism of C₄ plants is generally saturated at the current ambient atmospheric CO₂ (Ziska & Bunce, 1997), although there is evidence for stimulation of photosynthesis at eCO₂ even in C₄ plants (Leakey et al., 2004; Long et al., 2004). Since there was no evidence of downregulation of photosynthesis in the maize or wheat plants at eCO₂, it is possible that photosynthetic demand for N was similar, or increased (i.e. in the wheat plants) relative to the ambient CO₂ plants, thus leading to little or no difference in tissue N concentration between the two CO₂ levels. This study therefore does not provide evidence of decreased nutritional value in maize or wheat as a result of downregulation of photosynthesis

at eCO₂. In the wheat plants growing in the MIX NPK soil treatment, eCO₂ significantly increased biomass accumulation. This is most likely because the stimulation of photosynthesis at eCO₂ in these plants interacted with the adequate supply of nutrient and water in this soil treatment. However, a significant increase in the C:N and C:P ratios was measured in these plants, providing some evidence of a dilution effect of these two elements (Taub & Wang, 2008). Although eCO₂, combined with high nutrient availability and adequate water supply, increased biomass accumulation and nutrient content in the MIX NPK wheat plants, this positive effect seemed to have been cancelled out by the dilution of some elements. In contrast to wheat, photosynthetic rate and biomass accumulation of maize plants were not affected at eCO₂ under well-watered conditions, results which are in agreement with the literature (Ainsworth & Rogers, 2007; Markelz, Strellner & Leakey, 2011). Additionally, no evidence of dilution of any of the elements was found in the C₄ crop. However, in drought conditions at eCO₂, maize plants are able to improve WUE as a result of decreased stomatal conductance and sustained intercellular [CO₂], which can subsequently lead to increased biomass accumulation (Allen et al., 2011). I therefore suggest that increased biomass accumulation in maize plants under drought conditions, combined with reduced mass flow delivery of nutrients at eCO₂, can potentially decrease tissue nutrient concentration in maize plants as a result of dilution effects. However, this effect was not seen in the maize plants in this study possibly because they were not water-stressed.

Conclusion

The results from the experiment strongly suggest that the well-watered status of the soils may have allowed for an increase in the importance of diffusive processes for nutrient delivery to the rhizosphere when the plants were exposed to eCO₂, thereby making up for the reductions

in mass flow. As such, no decrease in tissue nutrient concentration was measured in the maize or wheat plants grown at eCO₂. These findings thus highlight the importance of considering soil water status when investigating and interpreting the effects of eCO₂-induced decreases in mass flow on the nutrient status of plants. Growth at eCO₂ under conditions of limited water availability, conditions which are not unlikely according to future climate predictions (Gerten et al., 2004; Fisher, Nakicenovic & Alfsen, 2007; Mu et al., 2007), may potentially negatively impact the quality and yield of important crops globally, as a result of decreased mass flow, and subsequently inadequate supply of nutrients.

List of figures

Figure 1: Variation in total biomass, root biomass and shoot-to-root ratio in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The boxes and horizontal lines represent the first and the third quartiles and the medians, respectively. The whiskers represent $1.5 \times$ the interquartile range and outliers above/below are shown as points. The diamonds represent the mean values and the filled circles indicate outliers. The F- and p-values of the main effects and interaction (when significant) from the analysis of variance done on the linear models fitted are displayed. Lower case letters indicate the significant differences between treatments as determined by Tukey's post-hoc comparisons.

Figure 2: Variation in average root diameter, total surface area and specific root length in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 3: Variation in stomatal conductance, transpiration rate, and photosynthetic rate in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 4: Variation in cumulative water loss measured by gravimetry in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 5: Variation in transpiration-driven mass flow delivery of N, P and K in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 6: Variation in transpiration-driven mass flow contribution of N, P and K to tissue content in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The dashed line represents 100% contribution to tissue content. Statistics as in Figure 1.


Figure 7: Variation in total tissue concentrations of N, P and K in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 8: Variation in total tissue contents of N, P and K in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Figure 1.

Figure 9: The relationship between transpiration-driven mass flow delivery of N, P and K, and total tissues contents of N, P and K, in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The intercept, slope, R² and p-values of the lines are displayed, as well as the p-value of the contrast between the 400 and 800 ppm line.

903 **Figure 10:** Variation in the C:N, C:P and C:K ratios in maize and wheat plants, grown in five
904 soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in
905 Figure 1.

906



908
909 **Figure 1**




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

Chapter 3

The effects of low vapour pressure deficit on the nutrient status of maize and wheat under well-watered conditions

Abstract

The effects of varying VPD levels on mass flow, biomass accumulation and nutrient status were investigated in maize and wheat plants. Plants were grown at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa in well-watered Soil and Sand. To demonstrate the importance of mass flow, plants grown in Sand were supplied nutrients in such a way that they had to rely exclusively on mass flow or diffusive processes (i.e. not interception) for nutrient delivery to their rhizosphere. Relative to the 1.613 kPa VPD level, cumulative water use (CW_L) was decreased on average by 40% and 17% at the 0.773 kPa level in the maize and wheat plants respectively, and by 48% and 35% at the 0.350 kPa level. In both soil treatments, mass flow contribution of most nutrients was in excess of tissue content in both species, even though a significant decrease was measured with decreasing VPD. Photosynthetic rates and stomatal conductance were little affected by VPD, but plants always accumulated more biomass at 0.773 kPa. Reduced CW_L may have impeded root-to-shoot transport of ions and reduced dry biomass accumulation in the maize and wheat plants at 0.350 kPa (-40% and -22% on average respectively, compared to 0.773 kPa plants). Tissue [NPK] was also lower (-13% [N], -41% [P] and -47% [K] respectively) in the 0.350 kPa VPD Sand-grown wheat plants, while increases in the proportion of finer roots may have alleviated reduced CW_L effects on tissue [NPK] in the C_4 species. Growth and nutrient concentrations in the maize and wheat plants were affected by their respective physiological and morphological response to varying VPD levels, rather than by changes in mass flow. The findings from this study suggest that negative effects of reduced mass flow may be alleviated under well-watered conditions.

Introduction

The difference in vapour pressure deficit (VPD) between leaf tissues and the atmosphere drives water evaporation from plants (Lihavainen et al., 2016). Low VPD decreases evaporative demand from the atmosphere, and as such, decreases plant transpiration rates, while the opposite is observed when plants are exposed to high VPD conditions (Adams & Hand, 1993; Yang et al., 2012). Varying VPD can have important effects on the growth and nutritional status of plants. An increase in air humidity decreases evaporative demand from the atmosphere and promotes stomatal opening, thereby facilitating CO₂ influx into the leaf (Bunce, 1984). Therefore, low VPD levels tend to increase biomass accumulation in plants (Leuschner, 2002; Lihavainen et al., 2016). However, the uptake of nutrients by the root systems of plants growing at low VPD may be reduced due to decreases in mass flow delivery of nutrients as a result of decreased plant transpiration (Barber, Walker & Vasey, 1963; Cramer, Hawkins & Verboom, 2009). Similarly, in a meta-analysis done by Mcgrath & Lobell (2013), a decline in nutrient concentrations due to reductions in transpiration has been reported in a variety of crops grown at elevated atmospheric [CO₂] ('eCO₂').

Transpiration-driven mass flow has been shown to be an important process affecting below-ground nutrient uptake in plants (Barber, 1995; Cramer, Hoffmann & Verboom, 2008). For example, Barber (1995) estimated that mass flow contributed to 79% N, 5% P, 18% K, 73% Ca, 88% Mg and 95% S in *Zea mays* plants growing in a fertile loam. Apart from air humidity and atmospheric [CO₂], plant transpiration can also be regulated by availability of below ground nutrients. In a study done by Cramer, Hoffmann & Verboom (2008), a first set of *Ehrharta calycina* plants had direct access to nutrients via interception, mass flow and diffusion, while a second set had to rely exclusively on mass flow or diffusive processes for nutrient delivery and uptake. As a result, the plants in the second set had a 63% higher transpiration rate and a 72% higher stomatal conductance, thus showing a nutritional regulation

of mass flow in this C₃ plant species. In a similar experimental setup, Rose et al. (2018) showed that C₄ grasses were unable to regulate their water flux in response to varying nutrient availability and accessibility. Thus, C₄ plants can potentially be more severely affected than C₃ plants in terms of their nutrition under conditions of reduced transpiration and low nutrient availabilities.

Several mechanisms, in addition to increased stomatal opening (Bunce, 1984; Gisleröd & Nelson, 1989; Lihavainen et al., 2016), have been proposed for the reported increases in biomass accumulation in C₃ plants under humid conditions. Such mechanisms include increases in leaf turgor and leaf water potential (Leuschner, 2002), which has been shown to increase leaf expansion in plants (Acevedo, Hsiao & Henderson, 1971; Kramer & Boyer, 1995). Altered root morphology is another mechanism that has been reported to lead to increases in biomass in plants growing at elevated air humidity. In a long-term study comparing the respective responses of silver birch and aspen trees to varying VPD levels, Rosenvald et al. (2014) initially found a decrease in growth in both species when grown at low VPD. However, while suppression of growth persisted in the aspen trees, the silver birch trees acclimatised to the high humidity environment and increased biomass accumulation by increasing the proportion of fine-root biomass and altering root morphology. This suggests that the response to increasing air humidity is not necessarily uniform amongst C₃ species.

In contrast, low humidity tends to decrease biomass accumulation in C₃ plants. For example, reductions in yield have been reported in wheat, kale, sugar beet (Ford & Thorne, 2016) and soya beans (Woodward & Begg, 1976) when exposed to high VPD levels. Reduced growth at low humidity may be explained by leaf water stress arising as a result of the rate of water uptake by the root systems being insufficient to meet the water demand of high transpiration rates (Hoffman, 1979; Ranathunge, Steudle & Lafitte, 2003; Kuwagata et al., 2012). Additionally, decreases in stomatal conductance to prevent excessive water loss under

conditions of high evaporative demand (Mott & Parkhurst, 1991; Oren et al., 1999; Sinclair, Zwieniecki & Holbrook, 2008) can lead to restriction of growth in C_3 plants. While transpiration and stomatal conductance tend to increase with increasing VPD, often these increases tend to reach a maximum at VPD levels above 2.0 kPa (Turner, Schulze & Gollan, 1985; Comstock & Ehleringer, 1993), which has been demonstrated to reduce growth in wheat as a result of reduced CO_2 influx (Condon et al., 2002). The stomatal and transpiration response of C_3 crops to increasing VPD levels can however be modulated by the water status of the soil in which the plants are growing. For example, Xue et al. (2004) experimentally demonstrated that transpiration rates in winter-wheat increased with increasing VPD under well-watered conditions, but decreased with increasing VPD under water-stressed conditions. Thus, C_3 plants exposed to high VPD levels, but with adequate water supply may show little to no decreases in their nutrient status as a result of unrestricted mass flow delivery of nutrients.

Since the photosynthetic apparatus of C_4 plants is usually saturated at ambient CO_2 (Ziska & Bunce, 1997), it is argued that decreased stomatal conductance at high VPD levels will have little to no effect on net photosynthetic CO_2 assimilation in these plants (Bunce, 1983; 1993). In a study examining the effects of eCO_2 on stomatal sensitivity to varying VPD levels, assimilation rate was not altered by varying VPD levels neither at ambient nor at eCO_2 in *Amaranthus hypochondriacus*, a C_4 species, while in *Helianthus annuus*, a C_3 species, insensitivity of stomatal conductance to changes in VPD was only found at eCO_2 (Bunce 1993). In another study, no net effect of low air humidity was found on photosynthesis and biomass accumulation in a C_4 plant, *Panicum maximum*, even though stomatal conductance was reduced (Kawamitsu, Yoda & Agata, 1993). In that same study, it was found that stomatal closure induced by low air humidity led to reductions in photosynthesis and carbon assimilation in *Oryza sativa*, a C_3 crop. Similarly, in the Negev desert of Israel, Lange et al. (1975) demonstrated that decreases in stomatal conductance led to greater decreases in net

photosynthesis at low VPD in C₃ plants, compared to C₄ plants, in which the reductions were minimal. This suggests that C₄ plants are potentially more resilient against reductions in growth than C₃ plants under low air humidity conditions. However, in a study comparing the effects of high VPD on photosynthesis and stomatal conductance in four C₄ species, Bunce (1982) showed that stomatal conductance and photosynthesis were decreased, in *Amaranthus hybridus*, *A. hypochondriacus* and *Zea mays*. However, in *Portulaca oleracea*, while stomatal conductance decreased, no decrease in net photosynthesis was observed. The author argued that increased mesophyll conductance, a nonstomatal limitation to photosynthesis, was responsible for the lack of change in photosynthesis in *P. oleracea* at high VPD. The above finding therefore suggests that the response to increasing VPD levels is not necessarily uniform amongst C₄ species. The photosynthetic response of C₄ species under humid conditions seem to however be similar to C₃ species. When comparing the effects of VPD on two C₄ species, *Amaranthus hybridus* and *Portulaca oleracea*, and a C₃ species *Chenopodium album*, Bunce (1983) found that net photosynthesis was increased in all three species when humidity levels were increased, which the author attributed to increased stomatal opening in all the species, leading to increased influx of CO₂. While there is evidence for the effects of varying VPD levels on the growth and development of C₄ plants (Bunce, 1983; Kawamitsu, Yoda & Agata, 1993; Yang et al., 2012), the effects of varying VPD levels on mass flow of nutrients has seldom been considered in these plants, even though changes in stomatal conductance and transpiration have been documented (Tanner & Beevers, 1990).

Increased biomass accumulation is often observed in plants growing at low VPD, since stomatal opening and increases in leaf expansion due to increases in leaf turgor pressure are favoured under humid conditions (Acevedo, Hsiao & Henderson, 1971; Kramer & Boyer, 1995; Leuschner, 2002; Lihavainen et al., 2016). However, decreases in transpiration at low VPD as a result of reduced evaporative demand can also negatively impact growth and nutrient

concentrations in plants. Tullus et al. (2012) measured a 30% decrease in xylem sap flow, a reduction in growth rate, as well as a significant decrease in leaf N content and concentrations in aspen trees growing in a high humidity environment. The authors attributed this decrease to an impairment of N uptake as a result of decreased mass flow. Additionally, Sellin et al. (2013) experimentally showed decreases in growth and in leaf concentrations of N and P in silver birch trees grown under elevated air humidity, also attributed to a decrease in water flux as a result of decreased transpiration rates. However, Tanner & Beevers (1990; 2001) experimentally demonstrated that a decrease in transpiration rates in C₃ and C₄ plants hydroponically grown under humid conditions did not affect growth or nutrient concentrations. Additionally, Leuschner (2002) did not find any correlations between transpiration rates and nutrient concentrations in eight herbaceous plants grown hydroponically at low VPD levels and attributed the decrease in nutrient concentrations to dilution effects as a result of increased growth. While Cramer, Hawkins & Verboom (2009) argue that transpiration has an important role in nutrient delivery to the rhizosphere of plants grown in soil, they argue it plays no role in terms of the uptake of minerals across the root membrane, which are instead achieved by transporters such as the Ammonium Transporter (AMT) or the Nitrate Transporter (NRT) for the uptake of NH₄⁺ or NO₃⁻ respectively (Crawford & Glass, 1998; Gojon et al., 2011; Xu, Fan & Miller, 2012). This potentially explains why a decrease in transpiration at low VPD does not alter the tissue nutrient status of plants grown hydroponically. However, in their experiment, Tanner & Beevers (1990) included a treatment of maize plants grown in soil and again found no effect of elevated air humidity (and consequent reduction in transpiration) on the nutrient status of these plants. Contrasting findings from the literature in terms of the effects of reduced transpiration on the nutrient status of plants grown under humid conditions therefore calls for a better understanding of the role and importance of mass flow delivery of nutrients.

In this study, I tested the effects of varying VPD levels on water use, biomass accumulation and nutrient status in maize and wheat plants. Experiments in which plants are grown at low VPD levels allow isolation of the effects of decreased transpiration on nutrient concentrations and delivery in plants. Although eCO₂ also leads to decreased transpiration rates in both C₃ and C₄ plants (see Chapter 2), it also has effects on other physiological process (i.e. photorespiration, Rubisco investment, root architecture and physiology), whereas VPD levels mainly affect transpiration rates. Thus, I expect decreasing VPD to affect biomass accumulation and nutrient status in a similar manner in wheat (C₃) and in maize (C₄). The experiment was performed under well-watered conditions so as to identify VPD effects on plant physiology independently of soil-water effects. Plants were grown at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa in well-watered Soil and Sand. To demonstrate the importance of mass flow, plants grown in Sand were supplied nutrients in such a way that they had to rely exclusively on mass flow or diffusive processes (i.e. not interception) for nutrient delivery to their rhizosphere. I first hypothesised that plant tissue nutrient concentrations and contents will decrease in both species with decreasing VPD as a result of reduced transpiration-driven mass flow of nutrients, and because of dilution effects due to increased biomass accumulation. I then hypothesised that root morphology will be altered in both species as a result of decreased delivery of nutrients by mass flow with increasing air humidity. Finally, I hypothesised that reduced mass flow at lower VPD levels will have a stronger effect on the nutrient status of plants that had to rely exclusively on mass flow or diffusive processes for nutrient uptake belowground (i.e. plants grown in the Sand treatment).

Methods

Plant germination and growth conditions

1110 Wheat (*Triticum aestivum* c. SST015, Kaap Agri, Malmesbury, 7300, South Africa) and maize
 1111 (*Zea mays* c. CAP 122-60, Capstone Seeds, Howick, 3290, South Africa) plants were grown
 1112 from seeds soaked in aerated water for 24 h and 48 h, respectively, before being sown in trays
 1113 of vermiculite. The trays were kept moist until the emerging seedlings were established. The
 1114 seedlings were transplanted into 0.20 m diameter pots, each respectively containing 2 kgs of
 1115 soil. 2 different soils were used in this experiment: a soil with high clay content (hereafter
 1116 referred to as ‘Soil’), and a sand with no clay content (hereafter referred to as ‘Sand’). The pots
 1117 containing Sand were equipped with a cylindrical central compartment to which plant roots did
 1118 not have access. These cylinders housed the fertiliser pellets so that plants growing in Sand had
 1119 to rely exclusively on transpiration-driven mass flow and diffusion so as to acquire nutrients
 1120 released from the pellets (Cramer, Hoffmann & Verboom, 2008). A hole was cut into the base
 1121 of each pot into which the central compartment was inserted and filled with sand. These
 1122 cylinders were made from PVC pipes (0.5 m diameter, 0.18 m long, 0.003 m wall thickness).
 1123 Holes of 0.0045 m in diameter were drilled into the cylinders, from top to bottom.. The holes
 1124 were covered with strips of 50 µm nylon mesh (Nylon 270, Ecotoa Enterprises, Kwadukuza,
 1125 South Africa) on the outer surface. Maize and wheat growing in Sand received a Long-Ashton
 1126 solution containing 4 mM NaNO₃, 0.67 mM NaH₂PO₄, 1.5 mM Na₂HPO₄, 2 mM K₂SO₄, 1.5
 1127 mM MgSO₄·7H₂O, 4 mM CaCl₂·2H₂O, 0.1388 mM H₃BO₃, 0.0208 mM MnSO₄·4H₂O, 0.0023
 1128 mM ZnSO₄·7H₂O, 0.0033 mM CuSO₄·5H₂O, 0.0002 mM Na₂MoO₄·2HO and 0.0899 Fe
 1129 EDTA (Hewitt, 1966) every third day during the first 9 days of growth. These plants were first
 1130 supplied with excess water before being supplied with 200 mL of the Long-Ashton solution.
 1131 After the establishment period, 25 g of Osmocote® Exact® Patterned Release Fertilizer (The
 1132 Scotts Company Ltd, c/o Bord na Móna, Main street, Newbridge, County Kildare, Dublin,
 1133 Ireland) pellets containing (in %) 15 N; 4 P; 7.5 K; 1.8 Mg; 0.02% B; 0.047 Cu; 0.40 Fe; 0.06
 1134 Mn; 0.020 Mo and 0.015 Zn were introduced into the central PVC compartment in such a way

so as to distributed them across the depth of the pot. Plants growing in Soil did not receive any fertiliser throughout the growth experiment as the amount of available nutrients in this soil was deemed adequate (see below) to sustain growth of maize and wheat plants throughout the experiment. These soil types and fertiliser treatments were replicated so as to have wheat and maize plants subject to them in high, intermediate and low VPD levels. The experiment was conducted in three growth rooms, set at either 55%, 75% or 95% RH. RH and temperature in the growth rooms were measured twice daily with a Brannan whirling hygrometer (1HYBR-137452, Prospectors Supplies Pty Ltd, Australia). The average RH and temperature during the growth experiment in the growth room set at 55% RH were 49.82% and 25.25 °C (corresponding VPD = 1.613 kPa), 76% and 25.28 °C (corresponding VPD = 0.773 kPa) in the 75% RH room, and 89.12% and 25.25 °C corresponding (VPD = 0.350 kPa) in the 95% RH room.

Field capacity and water use

Prior to running the experiment, the respective field capacities of the two soil types were determined. This was done by supplying excess water to pots containing 200 g of Soil or Sand that had been oven-dried 7 d at 70°C to remove any moisture. The pots were covered with plastic sheets to prevent evaporation and were allowed to settle for 24 h. Field capacity of the different soil types was then determined as the dry weight percentage of the difference between the dry weight of the soil and the wet weight after 48 h after they had settled (Kirkham, 2005). The field capacities were 26 and 12% for the Soil and the Sand respectively. Throughout the experiment, the soils were kept between 70-100% of their respective field capacities by gravimetric measurement of water content. These field capacities were chosen to allow unrestricted mass flow (Marschner & Rengel, 2011). Water use of plants and evaporation rates differed between the VPD treatments. Typically, in high VPD conditions, plants were watered daily, every two days in the intermediate conditions, and every 3 days in the low VPD

conditions. For each soil type, 7 additional control pots filled with the same amount of soil (2 kg) were randomly distributed in the three growth rooms set at either 55%, 75% or 95% RH. These control pots were subject to the same watering regimes as the pots with the plants. This was done so as to estimate the cumulative water use per plant throughout the growth experiment by gravimetry. Guttation was observed every morning at the tip of the leaves on most maize plants growing in Soil at the low VPD level during the last two weeks of the experiment.

Soil nutrient availability in Soil

A soil column was constructed from a 20-mL syringe with the plunger removed, and a disc of Whatman No. 1 filter paper was placed at the bottom. The filter paper was topped with glass wool to prevent loss of sediment. Oven-dried soil was filled to the 20 mL mark and water added to field capacity. The soil column was inserted through a hole in the lid of a 50-mL Falcon centrifuge tube, placed in the tube, and centrifuged (Eppendorf Centrifuge 5810R) at 150 g for 2 min. The volume of water collected in the centrifuge tube was taken as a relative measure of the mass flow displaceable soil solution. The mass flow technique using low-speed centrifugation was a slight modification of the gravity method described by (Barber, 1995a). The elemental determination in the water extracts were performed by the Elsenburg Laboratory (Western Cape Department of Agriculture, Stellenbosch, South Africa, 33.845259 S; 18.834722 E). The concentrations of P, K, Ca, Cu, Fe, Mn and Zn were determined directly using a Radial ICP Spectrometer ICAP 7600 (Thermo Fisher Scientific, Sunnyvale, CA, USA), while N was determined on a BR-71394 DA Gallery Analyzer (Thermo Fisher Scientific, Sunnyvale, CA, USA). The water extracts from the Soil contained (in mg l⁻¹) 112.3 N, 23.4 P, 290 K, 291 Ca, 0.038 Cu, 0.01 Mn, 0.3 Fe and 0.126 Zn.

Gas exchange measurements

1183 All plant gas exchange measurements were conducted on fully expanded leaves using a LI-
1184 6400 (Li-Cor, Lincoln, NE, USA) Portable Photosynthesis System equipped with 2 x 3 mm
1185 broadleaf cuvette and an integrated red/blue LI-6400-02B LED light source (Li-Cor, Lincoln,
1186 NE, USA), on well-watered plants. I set the RH in the cuvette to either 55%, 75% or 95%
1187 depending on the RH level to which the plants being measured were exposed. Reference CO₂
1188 was set to 400 ppm and block temperature was set to 25°C, to match the temperature at which
1189 the plants were growing at. The variables measured were photosynthetic rate, stomatal
1190 conductance, internal leaf carbon and transpiration rate. Water use efficiency (WUE) was then
1191 calculated as the ratio of photosynthetic rate to transpiration rate.

1192 *Plant harvest*

1193 The wheat and maize plants were harvested after 4 weeks of growth. The soil was carefully
1194 washed away with water on a 2 mm sieve to minimise root loss. The plants were separated into
1195 root and shoot, dried with paper towel, and immediately weighed to obtain fresh weight. The
1196 leaves were then scanned with an LI-3100 Area Meter (Li-Cor, Lincoln, NE, USA) so as to
1197 obtain the leaf area and the specific leaf area (SLA) of each plant. The harvested plants were
1198 then oven-dried at 70°C for 48 h, after which dry weight was determined. The difference
1199 between foliar fresh weight and foliar dry weight was divided by the foliar fresh weight to
1200 obtain foliar water concentration. A representative sub-sample that constituted at least 15% of
1201 the total fresh weight, and that included the whole length of the root system of each plant was
1202 carefully cut off (Vandamme et al., 2013) and used to determine average root diameter (mm)
1203 and total root surface area (m²) using a STD4800 scanner and WinRHIZO version 2013a
1204 (Regent Instruments, Canada). Specific root length (SRL, m g⁻¹) was calculated as the total root
1205 length (m) divided by the root dry weight (g). The root characteristics were converted to whole-
1206 root results by multiplying by their respective conversion factors to 100% of the root.

1207 *Mass flow delivery and contribution*

1208 Mass flow delivery of nutrients was calculated as the cumulative amount of water transpired
1209 by a plant throughout the growth experiment multiplied by the concentration of that nutrient in
1210 the soil solution. Since delivery of a nutrient does not necessarily equate to acquisition, mass
1211 flow contribution to tissue concentration of that nutrient was then calculated as a percentage of
1212 the total concentration of that nutrient in the plant. The remaining nutrient content of the plants
1213 was assumed to have been acquired by diffusion and/or root interception (Barber, 1995).

1214 *Foliar elemental analyses*

1215 The dried foliar and root samples were ground to a fine powder using a grinder or a ball-mill
1216 (MM200, Retsch, Germany). The powdered samples were placed in Perspex rings sealed with
1217 4 µm Polypropylene thin film (Chemplex Industries Inc, Florida, USA) and introduced to a
1218 SPECTRO XEPOS XRF spectrometer (SPECTRO, AMATEK materials analysis division,
1219 Kleve, Germany). Analyses were conducted using the X-LabPro 5 software, which
1220 incorporates the universal ‘Turbo Quant Powders’ method. The instrument was calibrated by
1221 using a certified standard GBW07312 (National Research Center for CRMs, Beijing, China),
1222 for which elemental concentrations were obtained from NOAA Technical memorandum NOS
1223 ORCA 68 (1992). Only the elements that were within the machine’s detection limits were
1224 included.

1225 The carbon isotope ratio ($\delta^{13}\text{C}$) and the total N concentration of the powdered foliar and
1226 root samples were determined using mass spectrometry. Samples were weighed into tin
1227 capsules (5 x 9 mm; Sántis Analytical, Teufen, Switzerland, with *ca.* 2 mg of foliar powder
1228 used for analysis). The samples were combusted in a Flash 2000 organic elemental analyser
1229 and the gases passed into a Delta V Plus isotope ratio mass spectrometer (IRMS) via a Conflo
1230 IV gas control unit (all from Thermo Scientific, Bremen, Germany). In-house standards and

one IAEA (International Atomic Energy Agency) standard (USGS25) were used to calibrate the results. Nitrogen concentration was expressed relative to atmospheric nitrogen (Evans, 2001).

Data analyses

All statistical analyses were performed using R Statistical Software (R Core Team, 2020). Comparisons of all the response variables were made using linear models with VPD level and soil type as explanatory variables, fitted as interactions, or as separate main effects if the interaction was not significant. Mass flow data was available only for plants growing in Soil, and thus comparisons were made using linear models with VPD levels as the explanatory variable, and mass flow delivery and mass flow contribution of nutrients respectively as response variables. Analyses of variance were done on all the fitted linear models to determine the significance of the main effects using the ‘car’ package (Fox & Weisberg, 2011). Where main effects significantly explained variation, pairwise comparisons (Tukey post-hoc tests) of group levels means were then performed using the package ‘emmeans’ (Lenth, 2018). All plots were generated using the ‘ggplot2’ package (Wickham, 2016).

Figures 1 to 8 (in Chapter 3) and Appendices 13 to 21 show the effects of soil treatment and VPD on morphological, physiological and nutrient variables in the maize and wheat plants. The ANOVA results indicating the significance of each factor are shown in each plot, with interaction terms excluded if they were not significant ($p > 0.05$). Note that a significant interaction term implies that the effect of VPD varied between soil treatments.

Results

Total biomass, root biomass, shoot-to-root ratio and root characteristics

Total biomass was higher at the intermediate VPD level in both maize and wheat plants, in both soil treatments (Fig. 1). Root biomass was higher at the intermediate VPD level in maize plants in both soil types, while there was no difference in root biomass between the VPD levels within each soil type in wheat plants (Fig. 1). Shoot-to-root ratios were highest in maize plants growing at the intermediate VPD level in Sand, with no difference measured between the VPD levels in Soil (Fig. 1). There was no difference in shoot-to-root ratio between the VPD levels within each soil type in wheat plants.

Average root diameter and total root surface area were higher in maize plants growing at the intermediate VPD level in both Sand and Soil, while no difference in these variables was measured between the VPD levels within each soil type in wheat plants (Fig. 2). There was no difference in specific root length between the VPD levels within each soil type in maize plants (Fig. 2). In wheat, specific root length was lowest at the intermediate VPD level in Sand, and highest at the high VPD level in Soil.

Gas exchange measurements, water use and carbon isotopic ratios

Neither VPD nor soil treatment affected stomatal conductance or photosynthetic rate in maize (Fig. 3). Photosynthetic rate was not affected by VPD in wheat plants either. In wheat plants, stomatal conductance was lowest at the intermediate VPD level in both soil types. Transpiration rate was highest at the high VPD level and lowest at the low VPD level in maize in both soil types (Fig. 3). Transpiration rate in wheat plants was highest at the high VPD level in both soil types, with no difference observed between the two other VPD levels. Cumulative water use per plant measured gravimetrically was highest at the high VPD level and lowest at the low VPD level in maize and wheat plants in both soil types (Fig. 4). Variation in foliar water concentrations could potentially indicate if any water stress arose as a result of high transpiration rates. Water concentration was lowest in maize growing at the intermediate VPD

level in Soil, and lowest at the high VPD level in Sand (Fig. 4). There were no differences in foliar water concentrations amongst the VPD levels in wheat plants. WUE was highest at the low VPD level and lowest at the high VPD level in maize in both soil types (Fig. 5). In wheat plants, WUE was higher at the low VPD level in both soil types, with no difference measured between the two other VPD levels. In maize, $\delta^{13}\text{C}$ was more negative at the high VPD level and less negative at the low VPD level in both Sand and Soil (Fig. 5). In wheat plants in Sand, $\delta^{13}\text{C}$ was less negative at the high VPD level and more negative at the low VPD level, while in Soil it was marginally more positive at the low VPD level. Variations in $\delta^{13}\text{C}$ at the different VPD levels can inform about potential stomatal limitations to CO_2 assimilation in both species (see Discussion).

Mass flow delivery of nutrients and contribution to tissue content in Soil

Mass flow delivered N, P, K, Ca, Cu, Mn, Fe and Zn was highest at the high VPD level and lowest at the low VPD level in both maize and wheat plants growing in Soil (Fig. 6, Appendices 13 and 14). Contribution of mass flow to tissue content of N, P, K, Ca, Cu, Mn, Fe and Zn in maize was the highest at the high VPD level, with no difference between the other VPD levels (Fig. 7, Appendices 15 and 16). In wheat plants, mass flow contributed N was highest at the high VPD level, while mass flow contributed Mn and Fe were lowest at the same VPD level. There was no difference in mass flow contributed P, K, Ca, Cu and Zn between the VPD levels in wheat plants. Contribution of mass flow to tissue content of N, P, K and Ca was above a 100% of tissue content at all VPD levels in maize and wheat plants growing in Soil (Fig. 6, Appendix 15). Additionally, mass flow contributed Cu, Mn, Fe and Zn were above 100% at all VPD levels in wheat plants growing in Soil (Appendices 15 and 16).

Tissue nutrient concentrations and contents

Tissue N, P, K concentrations were decreased at the high VPD level in both maize and wheat plants growing in Sand (Fig. 8). The concentrations of these nutrients were also decreased at the low VPD level in Sand grown wheat plants, but not in maize plants. Ca and Zn concentrations were the lowest Sand maize plants growing at the intermediate VPD levels, with little to no difference measured in Cu, Mn and Fe concentrations (Appendices 17 & 18). Little to no variation was measured in concentrations of Ca, Cu, Mn, Fe and Zn concentrations in Sand wheat plants. Concentrations of all the nutrients were lowest in Soil maize plants growing at the intermediate VPD level. In Soil wheat plants, concentrations of N, P, K and Ca did not vary between the VPD levels, while Cu, Mn and Fe concentrations were the highest at the high VPD level, and little to no difference was measured in Zn concentrations in these plants. The variation in the tissue content of all the nutrients in both maize and wheat plants, in both Sand and Soil, followed very similar patterns to total biomass at the different VPD levels (Appendices 19-21).

Discussion

As expected, transpiration rates and cumulative water use decreased with decreasing VPD in both maize and wheat, in both soil treatments. These decreases were mainly due to decreased evaporative demand, since stomatal conductance remained relatively unaffected by VPD in maize and was lowest at the intermediate VPD level in wheat. Additionally, VPD did not affect photosynthetic rates in either species. WUE increased with decreasing VPD, as a result of decreased transpiration rates. Since changes were nonetheless measured in biomass accumulation and nutrient concentrations in both species at the different VPD levels, the lack of variation in photosynthetic rate and stomatal conductance suggests that other factors such as

mass flow, soil water status, soil nutrient availability and accessibility, or dilution effects, influenced the growth and development of the plants at the different VPD levels.

Mass flow data was only available for the plants growing in the Soil. However, mass flow delivery of nutrients was likely decreased in a similar manner in Sand plants, since mass flow is calculated directly from cumulative water use, and cumulative water use was decreased in a similar manner in both soil treatments. Thus, reductions in mass flow, in both Soil and Sand, could potentially explain the reductions in biomass accumulation in the maize and wheat plants grown at the lowest VPD level. For example, in aspen and silver birch trees grown at low VPD, it has experimentally been demonstrated that decreases in mass flow, as a result of decreased transpiration, led to reductions in growth and foliar N content (Tullus et al., 2012; Sellin et al., 2015). In the present study however, mass flow contribution of N, P, K and Ca (as well Cu, Mn, Zn and Fe in wheat plants) was in excess of tissue contents of the maize and wheat plants, even at the lowest VPD level. The well-watered conditions (between 70-100% field capacity) to which the maize and wheat plants were subject at all the VPD levels and in both soil treatments may have allowed for excess delivery of these nutrients to the rhizosphere by mass flow, due to constant water flow, even though cumulative water use was reduced in the low VPD plants. Therefore, reductions in mass flow cannot fully explain the reductions in growth in the plants grown at this VPD level. One explanation could be that the low transpiration rates measured in both species at the low VPD level led to a reduction in xylem water movement, thus impeding root-to-shoot transport of ions within the plants. While Tanner & Beevers (1990; 2001) suggest that the quantity of transpiration-driven water flux in the xylem of plants considerably exceeds that required for movement of long-distance transport of ions, they do concede that this movement is usually accelerated by higher transpiration rates. For instance, transpiration mostly affects translocation rather than uptake of nutrients by roots in plants, but increased transpiration can nonetheless lead to increased ion uptake by increasing

the flux of elements through the roots' apoplastic pathway, potentially favouring greater uptake into the symplastic pathway (Marschner & Marschner, 2012). While mass flow delivery of nutrient to the rhizosphere does not equate to nutrient acquisition, the reduced transpiration rates may have decreased the nutrient uptake potential of the roots in the maize and wheat plants at the low VPD level, subsequently leading to reductions in biomass.

Tissue nutrient concentrations were, however, affected differently in the maize and wheat plants at the low VPD level. This contrast could be due to the decrease in root diameter at the low VPD level in the C₄ species, while no change in root diameter was measured in wheat plants (neither in Sand, nor in Soil) between the VPD levels. Increases in the proportion of fine roots have been demonstrated to be a response to low N or P availability (Gillespie, 1989; De Groot et al., 2003; Chapman et al., 2012; Luo et al., 2013) under conditions of varying VPD (Parts et al., 2013; Rosenvald et al., 2014). However, the ability for a plant to respond in terms of root morphology under conditions of low N or P availability and changing air humidity can differ between species, as well as between genotypes (Chapagain, 2014; Rosenvald et al., 2014). Thus, in the present study, the decrease in root diameter in the maize plants growing in Sand at the low VPD level may have allowed them to maintain adequate tissue concentrations of P and K compared to their wheat counterparts, even though transpiration rates and cumulative water use were decreased to a similar extent in both species. It has indeed been demonstrated that a morphological adaptation to low shoot P and low-P stress was an increase in the proportion of fine-root biomass in species such as soybean, sunflower or maize (Fernandez & Rubio, 2015; Wen et al., 2017). Furthermore, tissue nutrients concentrations in the wheat plants seem to have been affected differently depending on whether they grew in the Soil or in Sand. It is possible that reductions in transpiration combined with the restriction on root interception for nutrient uptake in the Sand treatment led to the lower P and K concentrations in the Sand wheat plants compared to the Soil wheat plants at the lowest VPD

level. Thus, there is good evidence to support the hypothesis that effects of reduced water flux with decreased VPD are exacerbated in Sand, where plants had to rely exclusively on mass and flow and diffusion.

In Soil maize plants, even though mass flow and biomass accumulation were reduced at the lowest VPD level, they maintained relatively high tissue concentrations of N, P, K, Ca and Zn compared to the ones growing at the intermediate VPD level. It is possible that guttation (in addition to increases in the proportion of finer roots), which was frequently observed on these plants during the growth experiment, may have facilitated the uptake and movement of nutrients within these plants even when transpiration rates were reduced (Tanner & Beevers, 1990; Slewinski, Meeley & Braun, 2009; Singh, 2016). Tanner & Beevers (1990) also observed guttation in maize plants growing under conditions where the air humidity was above 95%. This process has been shown to happen frequently in this C₄ crop (Erhardt, 1978) and potentially demonstrates its importance with regards to nutrient uptake and transport from the root to the shoot in plants.

In both soil treatments, maize and wheat plants growing at the high VPD level also experienced a reduction in biomass accumulation. This result is in agreement with previous studies where growth at low relative humidity was found to be reduced in various crop species (Woodward & Begg, 1976; Ranathunge, Steudle & Lafitte, 2003; Kuwagata et al., 2012; Ford & Thorne, 2016). The reduction of growth at low humidity in these studies was attributed to leaf water stress arising as a result of the rate of water uptake by the root systems being insufficient to meet the high transpiration rates. Significantly lower foliar water concentrations in the maize plants growing in Sand at the highest VPD level could potentially indicate that these plants were water-stressed as a result of high transpiration rates, thus leading to lower biomass. However, the maize plants growing in Soil, and the wheat plants in both soil treatments at the highest VPD level, did not have significantly lower foliar water

concentrations, suggesting that they were not water stressed. There were little to no changes in the shoot-to-root ratios between the VPD levels in both the C₃ and C₄ plant species, potentially suggesting that adequate root systems may have allowed the high VPD plants to avoid getting water-stressed, in addition to the well-watered status of the substrates. Thus, leaf water stress due to high transpiration rates cannot explain the reduction in growth in most of the plants grown at low air humidity in the present study. Furthermore, transpiration-driven mass flow of nutrients was highest in the high VPD plants, due to their high transpiration rates and cumulative water use. Why then is a reduction in biomass accumulation still observed in most of the plants grown at high VPD?

I suggest that stomatal and non-stomatal limitations to carbon assimilation may have led to the reductions in biomass observed in both species at the high VPD level. Gas exchange measurements did not reveal any differences in photosynthetic rates or stomatal conductance potentially because these measurements were always conducted after the plants had been watered, and thus any VPD-induced fluctuations in these two variables may have been masked. $\delta^{13}\text{C}$, on the other hand, is an integrated measure over the whole growth period of the plants and may be more indicative of any limitations to carbon assimilation throughout their development. The relationship between WUE and $\delta^{13}\text{C}$ is well-established in the literature (Farquhar et al., 1989; Condon, Richards & Farquhar, 1993; Wright, Rao & Farquhar, 1994; Brück, Payne & Sattelmacher, 2000). The opposite trends in $\delta^{13}\text{C}$ observed with increasing VPD (thus, with decreasing WUE) respectively in maize and wheat, are in agreement with the study done by Madhavan, Treichel & O'Leary (1991), with the differing discrimination of ^{13}C by Rubisco and PEP-Carboxylase in the C₃ and C₄ plants respectively explaining these trends (Farquhar, O'Leary & Berry, 1982; Winter et al., 1982). The respective $\delta^{13}\text{C}$ trends in the C₃ and the C₄ species potentially indicate that stomatal opening was favoured at low VPD, while closure was favoured at high VPD (Morison & Gifford, 1983; Bowman et al., 1989). Thus,

even though the gas exchange measurements did not detect differences in stomatal conductance, partial stomatal closure at the high VPD level (inferred from the $\delta^{13}\text{C}$ trends) may have restricted the influx of CO_2 , potentially restricting carbon assimilation (Cornic, 2000; Chaves et al., 2002; Cernusak et al., 2019), thereby leading to decreases in biomass. Higher evaporative demand at high VPD nonetheless still leads to higher transpiration rate for a given stomatal aperture (Sanchez-Diaz et al., 2002), and may explain the highest measured transpiration rates and cumulative water use of the high VPD plants, even if there was partial stomatal closure. Apart from stomatal effects, Sharkey (1984) has experimentally demonstrated that high transpiration rates could lead to non-stomatal limitations to carbon assimilation as a result of reduced mesophyll capacity for photosynthesis, attributed to water potential drops within the mesophyll. Consequently, decreases in biomass accumulation due to stomatal and non-stomatal limitations in the maize and wheat plants grown at high VPD may explain lower concentrations of N, P and K in their tissue, as a result of lower demand for growth.

The higher biomass accumulation in both maize and wheat plants at the intermediate VPD level is likely due to the absence of the different limitations on growth imposed by the more extreme VPD level. However, in Soil-grown maize at the intermediate VPD level, a possible dilution effect of most of the nutrients is observed, as shown by their lowest concentrations. Similarly, Leuschner (2002) experimentally demonstrated that shoot nutrient concentrations in eight plants grown at low VPD levels decreased due to dilution effects as a result of carbon assimilation in plants grown at low VPD levels. In the present study, transpiration rate and cumulative water use of the maize and wheat plants grown at the intermediate VPD level were nonetheless significantly decreased relative to the plants growing at the highest VPD level. Thus, increased biomass accumulation, coupled with a reduction in transpiration, may have led to inadequate supply and movement of nutrients by mass flow, thereby leading to a dilution effect. This is especially true for nutrients such as Cu, Mn, Fe and

Zn which were not supplied in excess of physiological requirements by mass flow in the intermediate VPD maize plants. The same explanation may also be applicable to describe the dilution of Ca and Zn in the intermediate VPD maize plants growing in Sand, since mass flow was the main mechanism for the delivery of these nutrients to the rhizosphere in this soil treatment. The above findings demonstrate the potentially important role of mass flow for nutrient delivery and on plant nutrient status.

Conclusion

The findings from this experiment corroborate those from the eCO₂ experiment (Chapter 2), in that well-watered conditions allowed for the delivery of most nutrients by mass flow in excess of physiological requirements, in both species, even when transpiration rates and cumulative water use were reduced at the intermediate and lowest VPD level. As such, decreases in mass flow had little to no effect on nutrient concentration or biomass accumulation in either species. Rather, the physiological and respective morphological responses to VPD of the maize and wheat plants led to changes in biomass accumulation and nutrient concentrations at the different VPD levels.

List of figures

Figure 1: Variation in total biomass, root biomass and shoot-to-root ratio in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. The boxes and horizontal lines represent the first and the third quartiles and the medians, respectively. The whiskers represent $1.5 \times$ the interquartile range and outliers above/below are shown as points. The diamonds represent the mean values and the filled circles indicate

outliers. The F- and p-values of the main effects and interaction (when significant/present) from the analysis of variance done on the linear models fitted are displayed. Lower case letters indicate the significant differences between treatments as determined by Tukey's post-hoc comparisons.

Figure 2: Variation in average root diameter, total surface area and specific root length in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Figure 1.

Figure 3: Variation in stomatal conductance, transpiration rate, and photosynthetic rate in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Figure 1.

Figure 4: Variation in cumulative water use and foliar water concentration in maize and wheat plants, grown in two Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Figure 1.

Figure 5: Variation in water-use efficiency and in $\delta^{13}\text{C}$ maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Figure 1.

Figure 6: Variation in transpiration-driven mass flow delivery of N, P and K in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Figure 1. Corresponding air relative humidity (in %) at 25°C shown on the x-axis.

Figure 7: Variation in transpiration-driven mass flow contribution of N, P and K to tissue content in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. The dashed line represents 100% contribution to tissue content. Statistics as in Figure 1. Corresponding air relative humidity (in %) at 25°C shown on the x-axis.

1494 **Figure 8:** Variation in total tissue concentrations of N, P and K in maize and wheat plants,
1495 grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics
1496 as in Figure 1.

1497

1498




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

Chapter 4

General discussion and synthesis

New perspectives?

Many studies have reported decreases in the nutritional value of a range of socio-economically important crops globally when grown under conditions of elevated atmospheric CO₂ (‘eCO₂’) (Loladze, 2002; Taub, Miller & Allen, 2008; Mcgrath & Lobell, 2013; Zhu et al., 2018). These decreases apply to a range of macronutrients such as N, P, K and Ca, as well as micronutrients such as Fe, Mn, Cu and Zn (e.g. in Fig. 1).




Figure 1: The effects of eCO₂ on wheat grain concentration of 12 different elements. Data are from a global meta-analysis on wheat (Broberg, Högy & Pleijel, 2017). The dashed horizontal line represents the effect on N for comparison. Figure obtained from Uddling et al. (2018).

Currently, it is estimated that 25% and 31% of World’s population are iron and zinc-deficient respectively (Ezzati et al., 2004). This global malnutrition problem could be further exacerbated, since no decrease in atmospheric [CO₂] is expected before the end of the century, even under the best mitigation scenarios (Fisher, Nakicenovic & Alfsen, 2007). Several

hypotheses have been proposed for the decrease in nutritional value of important crops at eCO₂, from reduced root development, dilution effects due to enhanced carbohydrate production, downregulation of photosynthesis and reduced transpiration-driven mass delivery of nutrients (Taub & Wang, 2008). In Chapters 2 and 3, mass flow delivery of nutrients was reduced in maize and wheat plants by growing them at eCO₂ (Chapter 2), and by decreasing vapour pressure deficits ('VPD') (Chapter 3), in order to determine the importance of this process on nutrient delivery and acquisition in these two species. Reduced mass flow at eCO₂ is generally accepted as a cause for decreased nutritional value in C₃ plants, but C₄ plants remain understudied in that context, possibly due to their apparent lack of response to eCO₂. The findings from Chapter 2 reveal that reductions in mass flow at eCO₂ occur in both the C₃ crop (wheat) and the C₄ crop (maize), as a result of decreased stomatal conductance and transpiration rates. However, no change in the tissue nutrient concentrations was observed in neither species and therefore the hypothesis that eCO₂-induced reductions in mass flow lead to decreased nutritional value wheat and maize was not supported. The lack of attributed to the well-watered status of the soil treatments (70-100% field capacity), which may have allowed for mass flow and diffuses processes to interchangeably transport nutrients to their root systems, even though mass flow was decreased. The findings from Chapter 3 corroborate the findings from Chapter 2, whereby decreases in mass flow delivery of nutrients with increasing relative air humidity did not affect tissue nutrient concentrations in either species under well-watered conditions. Combining the findings from both chapters and from the literature, I first argue that there is not enough evidence to support the hypothesis that reductions in mass flow lead to decreased nutrient concentrations at the leaf level in wheat and maize plants under well-watered conditions. I then argue against the idea that the nutrient status of C₄ crops are generally less at risk of deleterious effects under conditions of elevated atmospheric [CO₂] (Myers et al., 2014; Uddling et al., 2018).

Importance of soil water status on effects of reduced mass flow at eCO₂ and low VPD

It could be argued that the lack of decrease in tissue nutrient concentrations in the maize and wheat plants grown at eCO₂ (Chapter 2) was due to ‘pot effects’. Potted experiments, such as the ones conducted in Chapters 2 and 3, can firstly lead to different water dynamics compared to field conditions, while small soil volumes may allow for unrestricted access to nutrients by the roots, regardless of whether mass flow is reduced or not (Liebering et al., 2004; Mcgrath & Lobell, 2013). However, in a recent meta-analysis that included data from free-air CO₂ (FACE) experiments as well as from chamber experiments, decreases of a range of nutrients was found in crop species such as wheat or rice grown at eCO₂ (Myers et al., 2014). This potentially suggests that ‘pot effects’ may not be responsible for the unchanged nutrient concentrations of the maize and wheat plants grown at eCO₂. It was instead suggested that the well-watered status of the soil treatments (kept between 70-100% field capacity) may have allowed for the adequate delivery of nutrients by mass flow to the rhizosphere, even when transpiration was decreased (i.e. at eCO₂). In their meta-analysis, Mcgrath & Lobell (2013) state that the importance of mass flow can vary depending soil water content and soil solution nutrient concentrations. Thus, in well-watered substrates, with high nutrient concentrations in the soil solution, a lesser amount of water displaced (e.g. by mass flow) may nonetheless supply adequate amounts of nutrients to the rhizosphere. I thus argue that when investigating the effects of reduced transpiration and mass flow on the nutrient concentrations in plants, it is important to consider how the nutrient and water status of the soil may mediate the importance of mass flow for delivery of nutrients. For example, in a study done by del Pozo et al. (2007), decreases in N concentration in wheat plants grown at eCO₂ were attributed to a decrease in mass flow of N, as a result of decreased transpiration. The wheat plants in this study were exposed to soil water conditions that replicated water availability in a rainfed agricultural system, in Salamanca, Spain (i.e. not watered to field capacity). The findings from the above

study thus demonstrates that reductions in mass flow under conditions of limited soil water availability may indeed lead to decreased nutrient concentrations.

Under well-watered conditions in the eCO₂ experiment (Chapter 2), mass flow contribution of most nutrients to tissue content in both the maize and wheat plants was nonetheless significantly decreased at eCO₂, and yet their nutrient status remained unchanged. In that experiment, mass flow delivery of nutrients was calculated as the cumulative water use of a plant throughout its lifetime, multiplied by the concentration of that element in soil solution (Barber, 1995). Oyewole, Inselsbacher & Näsholm (2014) argue that challenges associated with this method are that they do not take into account spatial and temporal variations in soil water content, as well as potential interactions between mass flow and diffusion. Diffusive processes happen due to concentration gradients in and about the rhizosphere as a result of active uptake by root of plants (Barber, 1995). Even when transpiration and mass flow are reduced, movement of nutrients to the rhizosphere may thus be facilitated by diffusion under well-watered conditions. The linear regressions between amount of nutrients delivered by mass flow and tissue nutrient contents in the eCO₂ experiment showed higher N, P and K contents at the intercepts (i.e. when 0 mg of N, P and K was delivered by mass flow) in both maize and wheat. This strongly suggests that diffusive processes may have compensated for reductions in mass flow at eCO₂. In agreement with these findings, Van Vuuren et al. (1997) found that diffusive processes compensated for reductions in mass flow at eCO₂ in wheat plants grown in wet conditions, allowing them to maintain higher foliar N, compared to the ones grown at eCO₂ under dry conditions. Thus, in the latter study, the negative effects of reductions in transpiration-driven mass flow at eCO₂ were alleviated under adequate water supply. McDonald et al. (2002) made a similar observation in *Populus deltoides* trees grown at eCO₂ under varying relative air humidity. In that study, it was stated that diffusive processes with

high soil moisture in the high humidity treatments compensated for the eCO₂-induced reductions in mass flow for the delivery of NO₃⁻ to the trees.

Since high water availability is a condition that is not always met under field conditions (Marschner & Rengel, 2011), especially in rainfed agricultural systems (Lobell et al., 2014), eCO₂-induced reductions in transpiration may severely impact the nutritional value of many important crops globally. Decreases in the nutrient concentrations of plants exposed to eCO₂ are often attributed to a range of different mechanisms (Taub & Wang, 2008; Mcgrath & Lobell, 2013; Myers et al., 2014), which seem to vary in importance depending on environmental factors, as well as on species. For example, under well-watered conditions, photosynthesis is more likely to be stimulated in C₃ plant species at eCO₂ (Ainsworth & Rogers, 2007), and this can potentially make them more prone to dilution effects compared to C₄ plants species. The results from Chapter 2 confirm this assertion, since it was found that photosynthesis was greatly stimulated by eCO₂ in wheat compared to the maize plants, under well-watered conditions. In the case of reductions in transpiration driven-mass flow, the results from the eCO₂ Chapter, combined with findings the literature, show that the effects of this particular mechanism at eCO₂ can be strongly dependent on factors such as soil water status, soil texture and soil nutrient availability (Van Vuuren et al., 1997; del Pozo et al., 2007; McDonald et al. 2002; Oliveira et al., 2010; Mcgrath & Lobell, 2013).

C₃ versus C₄ crops in the context of mass flow reductions

Many studies have argued that C₄ plants are little affected by eCO₂ (Wong, 1979; Reeves et al., 1994; Uddling et al., 2018) due to their photosynthetic mechanism usually being saturated at ambient CO₂ (Ziska & Bunce, 1997). However, it has been shown that growth of C₄ plants can be stimulated under drought conditions at eCO₂ due to improved water-use efficiency as a result of decreased stomatal conductance and sustained intercellular [CO₂] (Ainsworth &

Rogers, 2007; Allen et al., 2011). Additionally, the results from Chapters 2 and 3 show that decreases in stomatal conductance and transpiration can occur in both the C₃ and the C₄ species. Thus, decreases in mass flow at eCO₂ and at low VPD occurred regardless of photosynthetic pathway, results which are in agreement with previous findings the literature (Wong, 1979; Tanner & Beevers, 1990; Loladze, 2002; Ainsworth & Long, 2005). Under conditions of limited soil water availability and eCO₂, reported increases in biomass accumulation (Ainsworth & Rogers, 2007), combined with reductions in stomatal conductance and transpiration (as shown by the results in Chapter 2), may in fact put C₄ crops equally at risk as C₃ crops in terms decreases in nutritional value under changing climatic conditions (e.g. eCO₂), as a result of decreased mass flow delivery of nutrients in addition to dilution effects. This is of particular concern because both wheat and maize are major sources of proteins and carbohydrates for a great proportion of the global population (Myers et al., 2014; Uddling et al., 2018) and continued increased in atmospheric [CO₂] may exacerbate the already existing global malnutrition problem.

While mass flow delivered nutrients in excess of physiological requirements in both maize and wheat at all CO₂ and VPD levels respectively in both experiments (attributed to well-watered conditions), the total amount of nutrients delivered by mass flow was nonetheless significantly reduced at eCO₂ and with decreasing VPD. Thus, under conditions of low soil water availability and nutrient availability, reductions in transpiration (i.e. at eCO₂ or low VPD) will most likely reduce the amount of nutrients available in the rhizosphere for uptake by the plants (Mortensen, 1987). In addition, while transpiration tends to generally affect delivery of nutrients, it has been suggested than uptake across the root membrane could also be reduced when transpiration is decreased due to decreased element flux through the root apoplast (Marschner & Marschner, 2012). Furthermore, movement of nutrients from the root to the shoot can also be impeded as a result of decreased xylem water movement when transpiration

is reduced, as shown by the decrease in foliar N in aspen trees grown under humid conditions (Tullus et al., 2012). The findings from both Chapters 2 and 3, in combination with findings from the literature, thus demonstrate the importance of taking into consideration plant transpiration-driven water fluxes both in the soil, and within the plants, in tandem with other metabolic and physiological changes (Taub & Wang, 2008), for the preservation of the nutritional value and yield of socio-economically important C₃ and C₄ crops under changing climatic conditions.

Moving forward

While nutrient or protein concentrations of the edible parts of the maize and wheat plants were not investigated at the different CO₂ or VPD levels in the present, it is well established that nutrients such as N are mostly translocated from the vegetative parts of these plants during grain filling (Papakosta & Gagianas, 1991; Mae, 1997; Ntanos & Koutroubas, 2002). For example, 40% of the N required for grain development in wheat was translocated from the leaves, with the other 60% coming from the stem, glumes and roots (Simpson, Lambers & Dalling, 1983). In maize, it has been shown that 56% of the total N translocated to the grain was from the aboveground vegetative parts (Beauchamp, Kannenberg & Hunter, 1976). Thus, studies investigating changes in foliar nutrients under conditions of reduced mass flow, such as the present one, can therefore inform agricultural management practices on how to maintain, mitigate, or improve the nutritional status of these two crops under changing climatic conditions.

Adequate water availability and increased fertiliser supply can potentially alleviate the negative effects of eCO₂ on the nutrient status of maize and wheat (Kimball, 2006; Bunce, 2014), but the use of these strategies is not always feasible, due to associated costs and environmental problems. Water is often one of the most limiting resource in cultivated systems

globally (Gerten et al., 2004; Mu et al., 2007). In addition, with the changing climate, global increases in temperature are expected, while the opposite is expected for precipitation in many regions where socio-economically important crops are grown (Giorgi et al., 2001; Kling et al., 2003; Weltzin et al., 2003), thereby increasing the risk of drought events. Furthermore, in many cultivated systems, increased application of fertiliser is not always possible due to cost constraints (Wallace & Knausenberger, 1997), while excessive fertiliser inputs may lead to a whole range of environmental problems, the most common one being eutrophication (Galloway et al., 2008).

Moreover, water being a limited resource from crop production, there has been a drive to develop more water-use efficient genotypes in the last decade (Fletcher, Sinclair & Allen, 2007; Gholipour et al., 2010; Jyostna Devi, Sinclair & Vadez, 2010; Yang et al., 2012), and it is possible that this may exacerbate the negative effects of eCO₂-induced reductions in mass flow on the nutrient status of these crops. It has been shown that decreased transpiration in water-use efficient cultivars of many important crops under conditions of eCO₂, high temperatures, or high VPD, increases soil moisture conservation in early growth stages, thereby leading to increased biomass accumulation as a result of more water being available in the late growing stages (Hunsaker et al., 2000; Leahey et al., 2006). Based on the findings from the study done by Van Vuuren et al. (1997), I argue that decreased transpiration and reduced soil water movement in water-use efficient genotypes may lead to inadequate supply of nutrients by mass flow to the root system of these crops, thereby leading to decreased tissue nutrient content, under conditions of low soil water availability and increasing global atmospheric [CO₂]. Additionally, increased biomass accumulation is often measured in limited water environments at eCO₂ in C₄ crops such as maize and sorghum (Reeves et al., 1994; Ainsworth & Rogers, 2007). Decreases in the nutritional value may be even further exacerbated, by a

1719 combination of reduced mass flow and dilution effects, in more water-use efficient cultivars of
1720 these C₄ crops.

1721 Moving forward, in addition to breeding water-use efficient cultivars, a possible way
1722 to mitigate the impacts of increasing atmospheric [CO₂] on the quality and yield of important
1723 crops globally can be to breed for cultivars of C₃ and C₄ crops that display reduced sensitivity
1724 to eCO₂ (Myers et al., 2014). For example, reduced stomatal closure at eCO₂ could be achieved
1725 by genetically modifying the CO₂ cellular sensing mechanism of plants. Such modifications
1726 include lowering abscisic acid (ABA) synthesis or modifying the expression patterns of
1727 carbonic anhydrases (CA) under conditions of eCO₂ (Engineer et al., 2016). Both ABA and
1728 CA function in mediating stomatal responses to CO₂ (Jones & Mansfield, 1970; Stimler, Berry
1729 & Yakir, 2012).

1730

References

- Acevedo, E., Hsiao, T.C. & Henderson, D.W. 1971. Immediate and Subsequent Growth Responses of Maize Leaves to Changes in Water Status. *Plant Physiology*. 8(5):631–636.
- Adams, P. & Hand, D.J. 1993. Effects of humidity and Ca level on dry-matter and Ca accumulation by leaves of cucumber (*Cucumis sativus* L.) . *Journal of Horticultural Science*. 68(5):767–774.
- Ainsworth, E.A. & Long, S.P. 2005. What have we learned from 15 years of free-air CO₂enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist*. 165(2):351–372.
- Ainsworth, E.A. & Rogers, A. 2007. The response of photosynthesis and stomatal conductance to rising [CO₂]: Mechanisms and environmental interactions. *Plant, Cell and Environment*. 30(3):258–270.
- Allen, L.H., Kakani, V.G., Vu, J.C.V. & Boote, K.J. 2011. Elevated CO₂ increases water use efficiency by sustaining photosynthesis of water-limited maize and sorghum. *Journal of Plant Physiology*. 168(16):1909–1918.
- Barber, S.A. 1995. *Soil nutrient bioavailability: a mechanistic approach*. John Wiley & Sons.
- Barber, S.A. & Cushman, J. 1981. Nitrogen uptake model for agronomic crops. In *Modeling waste water renovation-land treatment*. New York, NY, USA: Wiley-Interscience. 382–409.
- Barber, S.A., Walker, J.M. & Vasey, E.H. 1963. Mechanisms for Movement of Plant Nutrients from Soil and Fertilizer to Plant Root. *Journal of Agricultural and Food*

1754 *Chemistry*. 11:204–207.

1755 BassiriRad, H., Reynolds, J.F., Virginia, R.A. & Brunelle, M.H. 1997. Growth and Root NO₃⁻
 1756 and PO₄³⁻ Uptake Capacity of Three Desert Species in Response to Atmospheric CO₂
 1757 Enrichment. *Australian Journal of Plant Physiology*. 24:353–358.

1758 BassiriRad, H., Gutschick, V.P. & Lussenhop, J. 2001. Root system adjustments: Regulation
 1759 of plant nutrient uptake and growth responses to elevated CO₂. *Oecologia*.
 1760 126(3):305–320.

1761 Baysdorfer, C. & Robinson, J.M. 1985. Sucrose and Starch Synthesis in Spinach Plants
 1762 Grown under Long and Short Photosynthetic Periods. *Plant Physiology*. 79(3):838–
 1763 842.

1764 Beauchamp, E.G., Kannenberg, L.W. & Hunter, R.B. 1976. Nitrogen Accumulation and
 1765 Translocation in Corn Genotypes Following Silking 1. *Agronomy Journal*.
 1766 68(2):418–422.

1767 Bernacchi, C.J., Kimball, B.A., Quarles, D.R., Long, S.P. & Ort, D.R. 2006. Decreases in
 1768 Stomatal Conductance of Soybean under Open-Air Elevation of [CO₂] Are Closely
 1769 Coupled with Decreases in Ecosystem Evapotranspiration. *Plant Physiology*.
 1770 143(1):134–144.

1771 Berntson, G. M.; Woodward, F.I. 1992. The Root System Architecture and Development of
 1772 *Senecio vulgaris* in Elevated CO₂ and Drought. *Functional Ecology*. 6(3):324–333.

1773 Berntson, G. 1994. Modeling Root Architecture - Are There Tradeoffs Between Efficiency
 1774 and Potential of Resource Acquisition. *New Phytologist*. 127(3):483–493.

1775 Berntson, G.M. 1944. Modelling Root Architecture : Are There Tradeoffs between Efficiency
 1776 and Potential of Modelling tradeoffs root between architecture : efficiency and are

- 1777 there potential of resource acquisition? *New Phytologist*. 127(3):483–493.
- 1778 Berntson, G.M. & Bazzaz, F.A. 1996. Belowground positive and negative feedbacks on CO₂
- 1779 growth enhancement. *Plant and Soil*. 187(2):119–131.
- 1780 Bloom, A.J., Caldwell, R.M., Finazzo, J., Warner, R.L. & Weissbart, J. 1989. Oxygen and
- 1781 Carbon Dioxide Fluxes from Barley Shoots Depend on Nitrate Assimilation. *Plant*
- 1782 *Physiology*. 91(1):352–356.
- 1783 Bloom, A.J., Smart, D.R., Nguyen, D.T. & Searles, P.S. 2002. Nitrogen assimilation and
- 1784 growth of wheat under elevated carbon dioxide. *Proceedings of the National Academy*
- 1785 *of Sciences*. 99(3):1730–1735.
- 1786 Bowes, G. 1993. Facing the Inevitable : Plants and Increasing Atmospheric CO₂. *Annual*
- 1787 *review of plant physiology and plant molecular biology*. 44:309–332.
- 1788 Bowman, W.D., Hubick, K.T., Caemmerer, S. Von & Farquhar, G.D. 1989. Short-Term
- 1789 Changes in Leaf Carbon in Salt- and Water-Stressed C₄ Grasses. 90(1):162–166.
- 1790 Broberg, M.C., Högy, P. & Pleijel, H. 2017. CO₂ -induced changes in wheat grain
- 1791 composition: Meta-analysis and response functions. *Agronomy*. 7(2):1–18.
- 1792 Brouder, S.M. & Volenec, J.J. 2008. Impact of climate change on crop nutrient and water use
- 1793 efficiencies. *Physiologia Plantarum*. 133(4):705–724.
- 1794 Brück, H., Payne, W.A. & Sattelmacher, B. 2000. Effects of phosphorus and water supply on
- 1795 yield, transpirational water-use efficiency, and carbon isotope discrimination of pearl
- 1796 millet. *Crop Science*. 40(1):120–125.
- 1797 Bunce, J.A. 1982. Low humidity effects on photosynthesis in single leaves of C₄ plants.
- 1798 *Oecologia*. 54(2):233–235.

- 1799 Bunce, J.A. 1983. Differential Sensitivity to Humidity of Daily Photosynthesis in the Field C₃
1800 and C₄ Species. *Oecologia*. 57(1):262–265.
- 1801 Bunce, J.A. 1984. Effects of humidity on photosynthesis. *Journal of Experimental Botany*.
1802 35(9):1245–1251.
- 1803 Bunce, J.A. 1993. Effects of doubled atmospheric carbon dioxide concentration on the
1804 responses of assimilation and conductance to humidity. *Plant, Cell and Environment*.
1805 16:189–197.
- 1806 Bunce, J.A. 2014. Corn Growth Response to Elevated CO₂ Varies with the Amount of
1807 Nitrogen Applied. *American Journal of Plant Sciences*. 05(03):306–312.
- 1808 Cakmak, I., Pfeiffer, W.H. & McClafferty, B. 2010. REVIEW: Biofortification of Durum
1809 Wheat with Zinc and Iron. *Cereal Chemistry*. 87(1):10–20.
- 1810 Cernusak, L.A., Goldsmith, G.R., Arend, M. & Siegwolf, R.T.W. 2019. Effect of vapor
1811 pressure deficit on gas exchange in wild-type and abscisic acid-insensitive plants.
1812 *Plant Physiology*. 181(4):157–1586.
- 1813 Chapagain, T. 2014. Root Architecture Variation in Wheat and Barley Cultivars. *American*
1814 *Journal of Experimental Agriculture*. 4(7):849–856.
- 1815 Chapin, F.S. 1980. The mineral nutrition of wild plants. *Annual Review of Ecology and*
1816 *Systematics*. 11:233-260
- 1817 Chapman, N., Miller, A.J., Lindsey, K. & Whalley, W.R. 2012. Roots, water, and nutrient
1818 acquisition: Let's get physical. *Trends in Plant Science*. 17(12):701–710.
- 1819 Chaves, M.M., Pereira, J.S., Maroco, J., Rodrigues, M.L., Ricardo, C.P.P., Osório, M.L.,
1820 Carvalho, I., Faria, T., Pinheiro, C. 2002. How plants cope with water stress in the
1821 field. Photosynthesis and growth. *Annals of Botany*. 89:907–916.

- 1822 Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J.
1823 Galloway, M. Heimann, C.
- 1824 Jones, C. Le Quéré, R.B. Myneni, S. Piao and P. Thornton 2013: Carbon and Other
1825 Biogeochemical Cycles. In: *Climate Change 2013: The Physical Science Basis.*
1826 *Contribution of Working Group I to the Fifth Assessment Report of the*
1827 *Intergovernmental Panel on Climate Change*. Stocker, T.F., D. Qin, G.-K. Plattner,
1828 M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley,
1829 Eds. Cambridge University Press, Cambridge, United Kingdom and New York, NY,
1830 USA
- 1831 Comas, L.H., Bouma, T.J. & Eissenstat, D.M. 2002. Linking root traits to potential growth
1832 rate in six temperate tree species. *Oecologia*. 132(1):34–43.
- 1833 Comstock, J. & Ehleringer, J. 1993. Stomatal response to humidity in common bean
1834 (*Phaseolus vulgaris*): Implications for maximum transpiration rate, water-use
1835 efficiency and productivity. *Australian Journal of Plant Physiology*. 20(6):669–691.
- 1836 Condon, A.G., Richards, R.A. & Farquhar, G.D. 1993. Relationships between carbon isotope
1837 discrimination, water use efficiency and transpiration efficiency for dryland wheat.
1838 *Australian Journal of Agricultural Research*. 44(8):1693–1711.
- 1839 Condon, A.G., Richards, R.A., Rebetzke, G.J. & Farquhar, G.D. 2002. Improving Intrinsic
1840 Water-Use Efficiency and Crop Yield. *Crop Science*. 42(February):122–131.
- 1841 Conroy, J. 1992. Influence of Elevated Atmospheric CO₂ Concentrations on Plant Nutrition.
1842 *Australian Journal of Botany*. 40(5):445.
- 1843 Cornic, G. 2000. Drought stress inhibits photosynthesis by decreasing stomatal aperture - not
1844 by affecting ATP synthesis. *Trends in Plant Science*. 5:187–188.

- 1845 Cousins, A.B., Adam, N.R., Wall, G.W., Kimball, B.A., Pinter, P.J., Ottman, M.J., Leavitt,
1846 S.W. & Webber, A.N. 2003. Development of C₄ photosynthesis in sorghum leaves
1847 grown under free-air CO₂ enrichment (FACE). *Journal of Experimental Botany*.
1848 54(389):1969–1975.
- 1849 Cramer, M.D. & Hawkins, H.J. 2009. A physiological mechanism for the formation of root
1850 casts. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 274(3–4):125–133.
- 1851 Cramer, M.D., Hoffmann, V. & Verboom, G.A. 2008. Nutrient availability moderates
1852 transpiration in *Ehrharta calycina*. *New Phytologist*. 179(4):1048–1057.
- 1853 Cramer, M.D., Hawkins, H.-J. & Verboom, G.A. 2009. The importance of nutritional
1854 regulation of plant water flux. *Oecologia*. 161(1):15–24.
- 1855 Crawford, N.M. & Glass, A.D.M. 1998. Molecular and physiological aspects of nitrate
1856 uptake in plants. *Trends in Plant Science*. 3(10):389–395.
- 1857 Cure, J.D. & Acock, B. 1986. Crop responses to carbon dioxide doubling: a literature survey.
1858 *Agricultural and Forest Meteorology*. 38(1–3):127–145.
- 1859 Davey, P.A., Parsons, A.J., Atkinson, L., Wadge, K. & Long, S.P. 1999. Does photosynthetic
1860 acclimation to elevated CO₂ increase photosynthetic nitrogen-use efficiency ? A study
1861 of three native UK grassland species in open-top chambers. *Environmental Research*.
1862 13(1):21–28.
- 1863 Drake, B.G., González-Meler, M.A. & Long, S.P. 1997. MORE EFFICIENT PLANTS: A
1864 Consequence of Rising Atmospheric CO₂? . *Annual Review of Plant Physiology and*
1865 *Plant Molecular Biology*. 48(1):609–639.
- 1866 Ehleringer, J.R., Cerling, T.E. & Helliker, B.R. 1997. C₄ photosynthesis, atmospheric CO₂ ,
1867 and climate. *Oecologia*. 112(3):285–299.

- 1868 Eng, R.Y.N., Tsujita, M.J. & Grodzinski, B. 1985. The effects of supplementary HPS lighting
1869 and carbon dioxide enrichment on the vegetative growth, nutritional status and
1870 flowering characteristics of *Chrysanthemum morifolium* Ramat. *Journal of*
1871 *Horticultural Science*. 60(3):389–395.
- 1872 Engineer, C., Hashimoto-Sugimoto, M., Negi, J., Israelsson-Nordstrom, M., Azoulay-
1873 Shemer, T., Rappel, W.-J., Iba, K. & Schroeder, J. 2016. CO₂ sensing and CO₂
1874 regulation of stomatal conductance: advances and open questions. *Trends in Plant*
1875 *Science*. 21(1):16–30.
- 1876 Erbs, M., Manderscheid, R., Hüther, L., Schenderlein, A., Wieser, H., Dänicke, S. & Weigel,
1877 H.J. 2014. Free-air CO₂ enrichment modifies maize quality only under drought stress.
1878 *Agronomy for Sustainable Development*. 35(1):203–212.
- 1879 Erhardt, J.F. 1978. Untersuchungen über Hydathoden. Technische Universität, München.
- 1880 Evans, L.T. 1993. *Crop Evolution, Adaptation, and Yield*. Cambridge University Press.
- 1881 Evans, R.D. 2001. Physiological mechanisms influencing plant nitrogen isotope composition.
1882 *Trends in Plant Science*. 6(3):121–126.
- 1883 Evans, J. & Seemann, J.R.. 1989. The allocation of protein nitrogen in the photosynthetic
1884 apparatus: Costs, consequences, and control. *Photosynthesis*. 183-205.
- 1885 Ezzati, M., Lopez, A.D., Rodgers, A. & Murray, C.J.L. 2004. Comparative quantification of
1886 health risks: global and regional burden of diseases attributable to selected major
1887 risks. *Comparative Quantification of Health Risks: Global and Regional Burden of*
1888 *Diseases Attributable to Selected Major Risk Factors*.
- 1889 Fangmeier, A., De Temmerman, L., Mortensen, L., Kemp, K., Burke, J., Mitchell, R., Van
1890 Oijen, M. & Weigel, H.J. 1999. Effects on nutrients and on grain quality in spring

1891 wheat crops grown under elevated CO₂ concentrations and stress conditions in the
1892 European, multiple-site experiment “ESPACE-wheat”. *European Journal of*
1893 *Agronomy*. 10(3–4):215–229.

1894 Fangmeier, A., Chrost, B. & Ho, P. 2000. CO₂ enrichment enhances flag leaf senescence in
1895 barley due to greater grain nitrogen sink capacity. *Environmental and Experimental*
1896 *Botany*. 44:151–164.

1897 Fangmeier, A., De Temmerman, L., Black, C., Persson, K. & Vorne, V. 2002. Effects of
1898 elevated CO₂ and/or ozone on nutrient concentrations and nutrient uptake of potatoes.
1899 *European Journal of Agronomy*. 17(4):353–368.

1900 FAO. (2013). Food and Agriculture Organization of the United Nations, FAOSTAT database.
1901 Rome: 703 Food and Agriculture Organization of the United Nations. URL
1902 <http://faostat.fao.org/>

1903 Farquhar, G.D., O’Leary, M.H. & Berry, J.A. 1982. On the relationship between carbon
1904 isotope discrimination and the intercellular carbon dioxide concentration in leaves.
1905 *Australian Journal of Plant Physiology*. 9(2):121–137.

1906 Farquhar, G.D., Hubick, K.T., Condon, A.G. & Richards, R.A. 1989. Carbon Isotope
1907 Fractionation and Plant Water-Use Efficiency. *Stable Isotopes in Ecological*
1908 *Research*. 21–40.

1909 Feng, Z., Rütting, T., Pleijel, H., Wallin, G., Reich, P.B., Kammann, C.I., Newton, P.C.,
1910 Kobayashi, K., Luo, Y. and Uddling, J. 2015. Constraints to nitrogen acquisition of
1911 terrestrial plants under elevated CO₂. *Global Change Biology*. 21:3152–3168.

1912 Fernandez, M.C. & Rubio, G. 2015. Root morphological traits related to phosphorus-uptake
1913 efficiency of soybean, sunflower, and maize. *Journal of Plant Nutrition and Soil*

- 1914 *Science*. 178(5):807–815.
- 1915 Fisher, B.S., Nakicenovic, N. & Alfsen, K. 2007. Issues related to mitigation in the long term
 1916 context. In *Climate change 2007: mitigation contribution of working group III to the*
 1917 *Fourth Assessment Report of the Inter-governmental Panel on Climate Change*. B.
 1918 Metz, O.R. Davidson, P.R. Bosch, R. Dave, & L.A. Meyer, Eds. Cambridge:
 1919 Cambridge University Press. 169–250.
- 1920 Fitter, A.H. 1987. An architectural approach to the comparative ecology of plant root
 1921 systems. *New Phytologist*. 106:61–77.
- 1922 Fletcher, A.L., Sinclair, T.R. & Allen, L.H. 2007. Transpiration responses to vapor pressure
 1923 deficit in well watered “slow-wilting” and commercial soybean. *Environmental and*
 1924 *Experimental Botany*. 61(2):145–151.
- 1925 Ford M.A., Thorne G.N. 1974 Effects of Atmospheric Humidity on Plant Growth. *Annals of*
 1926 *Botany*. 38:441–452.
- 1927 Fox, J. & Weisberg, S. 2011. *An {R} Companion to Applied Regression, Second Edition*.
 1928 Thousand Oaks CA: Sage.
- 1929 Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R.,
 1930 Martinelli, L.A., Seitzinger, S.P., Sutton, M.A. 2008. Transformation of the Nitrogen
 1931 Cycle : *Science*. 320:889–892.
- 1932 Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W. & Sitch, S. 2004. Terrestrial vegetation
 1933 and water balance - Hydrological evaluation of a dynamic global vegetation model.
 1934 *Journal of Hydrology*. 286(1–4):249–270.
- 1935 Ghannoum, O., Caemmerer, S. Von, Ziska, L.H. & Conroy, J.P. 2000. The growth response
 1936 of C₄ plants to rising atmospheric CO₂ partial pressure: a reassessment. *Plant, Cell*

- 1937 *and Environment*. 23(9):931–942.
- 1938 Gholipoor, M., Prasad, P.V.V., Mutava, R.N. & Sinclair, T.R. 2010. Genetic variability of
 1939 transpiration response to vapor pressure deficit among sorghum genotypes. *Field*
 1940 *Crops Research*. 119(1):85–90.
- 1941 Gifford, R.M., Barrett, D.J. & Lutze, J.L. 2000. The effects of elevated [CO₂] on the C:N and
 1942 C:P mass ratios of plant tissues. *Plant and Soil*. 224:1–14.
- 1943 Gillespie, A.R. 1989. Modelling nutrient flux and interspecies root competition in
 1944 agroforestry interplantings. *Agroforestry Systems*. 8(3):257–265.
- 1945 Giorgi, F., Whetton, P.H., Jones, R.G., Christensen, J.H., Mearns, L.O., Hewitson, B., Von
 1946 Storch, H., Francisco, R. & Jack, C. 2001. Emerging patterns of simulated regional
 1947 climatic changes for the 21st century due to anthropogenic forcings. *Geophysical*
 1948 *Research Letters*. 28. 3317–3320.
- 1949 Gisleröd, H.R. & Nelson, P. V. 1989. Effects of Humidity on Photosynthesis. *Scientia*
 1950 *Horticulturae*. 38:305–313.
- 1951 Gojon, A., Krouk, G., Perrine-Walker, F. & Laugier, E. 2011. Nitrate transceptor(s) in plants.
 1952 *Journal of Experimental Botany*. 62(7):2299–2308.
- 1953 De Groot, C.C., Marcelis, L.F.M., Van Den Boogaard, R., Kaiser, W.M. & Lambers, H.
 1954 2003. Interaction of nitrogen and phosphorus nutrition in determining growth. *Plant*
 1955 *and Soil*. 248(1–2):257–268.
- 1956 Hewitt, E. 1966. *Sand and water culture methods used in the study of plant nutrition*. 2nd ed.
 1957 Farnham Royal, England: Common Agricultural Bureau.
- 1958 Hocking, P.J. & Meyer, C.P. 1991. Carbon dioxide enrichment decreases critical nitrate and
 1959 nitrogen concentrations in wheat. *Journal of Plant Nutrition*. 14(6):571–584.

- 1960 Hoffman, G.J. 1979. Humidity. In *Controlled environment guidelines for plant research*.
1961 T.W. Tibbits & T.T. Kozlowski, Eds. London: Academic Press.
- 1962 Houshmandfar, A., Fitzgerald, G.J., O'Leary, G., Tausz-Posch, S., Fletcher, A. & Tausz, M.
1963 2018. The relationship between transpiration and nutrient uptake in wheat changes
1964 under elevated atmospheric CO₂. *Physiologia Plantarum*. 163(4): 516–529.
- 1965 Hunsaker, D.J., Kimball, B.A., Pinter, P.J., Wall, G.W., Lamorte, R.L., Adamsen, F.J.,
1966 Leavitt, S.W., Thompson, T.L., Matthias, A.D., Brooks, T.J. 2000. CO₂ enrichment
1967 and soil nitrogen effects on wheat evapotranspiration and water use efficiency.
1968 *Agricultural and Forest Meteorology*. 104(2):85–105.
- 1969 IPCC. 2014. *Climate change 2014: synthesis report.* ' *Contribution of Working Groups I, II*
1970 *and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
1971 *Change. Core Writing Team, RK Pachauri, LA Meyer Eds. IPCC, Geneva,*
1972 *Switzerland, 151*
- 1973 Jablonski, L.M., Wang, X. & Curtis, P.S. 2002. Plant reproduction under elevated CO₂
1974 conditions: A meta-analysis of reports on 79 crop and wild species. *New Phytologist*.
1975 156(1):9–26.
- 1976 Jones, R.J. & Mansfield, T.A. 1970. Suppression of stomatal opening in leaves treated with
1977 abscisic acid. *Journal of Experimental Botany*. 21(3):714–719.
- 1978 Jyostna Devi, M., Sinclair, T.R. & Vadez, V. 2010. Genotypic variation in peanut for
1979 transpiration response to vapor pressure deficit. *Crop Science*. 50(1):191–196.
- 1980 Kawamitsu, Y., Yoda, S. & Agata, W. 1993. Humidity Pretreatment Affects the Responses of
1981 Stomata and CO₂ Assimilation to Vapor Pressure Difference in C₃ and C₄ Plants.
1982 *Plant Cell*. 34(1):113–119.

- 1983 Kimball, B.A. 2006. The Effects of Free-Air [CO₂] Enrichment of Cotton, Wheat, and
1984 Sorghum. *Managed Ecosystems and CO₂*. 187:47–70.
- 1985 Kimball, B.A. 2016. Crop responses to elevated CO₂ and interactions with H₂O, N, and
1986 temperature. *Current Opinion in Plant Biology*. 31:36–43.
- 1987 Kimball, B.A. 2006. The effects of free-air [CO₂] enrichment of cotton, wheat and sorghum.
1988 In *Managed Ecosystems and CO₂. Case Studies, Processes and Perspectives*. S.P.J.
1989 Nösberger, S.P. Long, R.J. Norby, M. Stitt, Eds. Heidelberg, Berlin: Springer-Verlag.
1990 47–70.
- 1991 Kirkham, M.B. 2005. Field Capacity, Wilting Point, Available Water, and the Non-Limiting
1992 Water Rang. In *Principles of Soil and Plant Water Relations*. M.B. Kirkham, Ed.
1993 Academic Press. 101–115.
- 1994 Kling, G.W., Hayhoe, K., Johnson, L.B., Magnuson, J.J., Polasky, S., Robinson, S.K., Shuter,
1995 B.J., Wander, M.M., Wubbles, D., Donald, Z. 2003. *Confronting Climate Change in
1996 the Great Lakes Region: Impacts on our Communities and Ecosystems*. Union of
1997 Concerned Scientists, Cambridge, Massachusetts, and Ecological Society of America,
1998 Washington, D.C.
- 1999 Körner, C. 2006. Plant CO₂ responses: An issue of definition, time and resource supply. *New
2000 Phytologist*. 172(3):393–411.
- 2001 Kramer, P.J. & Boyer, J.N. 1995. *Water Relations of Plants and Soils*. San Diego: Academic
2002 Press.
- 2003 Kuehny, J.S., Peet, M.M., Nelson, P. V. & Willits, D.H. 1991. Nutrient Dilution by Starch in
2004 CO₂-enriched *Chrysanthemum*. *Journal of Experimental Botany*. 42(6):711–716.
- 2005 Kuwagata, T., Ishikawa-Sakurai, J., Hayashi, H., Nagasuga, K., Fukushi, K., Ahamed, A.,

- 2006 Takasugi, K., Katsuhara, M., Murai-Hatano, M. 2012. Influence of low air humidity
2007 and low root temperature on water uptake, growth and aquaporin expression in rice
2008 plants. *Plant and Cell Physiology*. 53(8):1418–1431.
- 2009 Lambers, H., Chapin, F.I. & Pons, T. 1998. *Plant physiological ecology*. New York, NY,
2010 USA: Springer-Verlag New York.
- 2011 Lange, O.L., Schulze, E.-D., Kappen, L., Buschbom, U. & Evenari, M. 1975. Photosynthesis
2012 of desert plants as influenced by internal and external factors. In *Perspectives in*
2013 *Biophysical Ecology. Ecological Studies v. 12*. D.M. Gates & R.D. Schmerl, Eds.
2014 Springer-Verlag, New York. 121–143.
- 2015 Larcher, W. 1995. *Physiological plant ecology*. Springer-Verlag, New York.
- 2016 Leakey, A.D.B. 2009. Rising atmospheric carbon dioxide concentration and the future of C₄
2017 crops for food and fuel. *Proceedings of the Royal Society*. 276(1666):2333–43.
- 2018 Leakey, A.D.B., Bernacchi, C.J., Dohleman, F.G., Ort, D.R. & Long, S.P. 2004. Will
2019 photosynthesis of maize (*Zea mays*) in the US Corn Belt increase in future [CO₂] rich
2020 atmospheres? An analysis of diurnal courses of CO₂ uptake under free-air
2021 concentration enrichment (FACE). *Global Change Biology*. 10(6):951–962.
- 2022 Leakey, A.D.B., Uribeharrea, M., Ainsworth, E.A., Naidu, S.L., Rogers, A., Ort, D.R. &
2023 Long, S.P. 2006. Photosynthesis, productivity, and yield of maize are not affected by
2024 open-air elevation of CO₂ concentration in the absence of drought. *Plant Physiology*.
2025 140(2):779–790.
- 2026 Leakey, A.D.B., Ainsworth, E.A., Bernacchi, C.J., Rogers, A., Long, S.P. & Ort, D.R. 2009.
2027 Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important
2028 lessons from FACE. *Journal of Experimental Botany*. 60(10):2859–2876.

2029 Russell Lenth (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. R
 2030 package version 1.4.1. <https://CRAN.R-project.org/package=emmeans>

2031 Leuschner, C. 2002. Air humidity as an ecological factor for woodland herbs: Leaf water
 2032 status, nutrient uptake, leaf anatomy, and productivity of eight species grown at low
 2033 or high vpd levels. *Flora*. 197(4):262–274. DOI: 10.1078/0367-2530-00040.

2034 Li, F., Kang, S. & Zhang, J. 2004. Interactive effects of elevated CO₂, nitrogen and drought
 2035 on leaf area, stomatal conductance, and evapotranspiration of wheat. *Agricultural*
 2036 *Water Management*. 67(3):221–233.

2037 Lieffering, M., Kim, H.Y., Kobayashi, K. & Okada, M. 2004. The impact of elevated CO₂ on
 2038 the elemental concentrations of field-grown rice grains. *Field Crops Research*. 88(2–
 2039 3):279–286.

2040 Lihavainen, J., Ahonen, V., Keski-Saari, S., Kontunen-Soppela, S., Oksanen, E. & Keinänen,
 2041 M. 2016. Low vapour pressure deficit affects nitrogen nutrition and foliar metabolites
 2042 in silver birch. *Journal of Experimental Botany*. 67(14):4353–4365.

2043 Lobell, D.B., Roberts, M.J., Schlenker, W., Braun, N., Little, B.B., Rejesus, R.M. &
 2044 Hammer, G.L. 2014. Greater sensitivity to drought accompanies maize yield increase
 2045 in the U.S. Midwest. *Science*. 344(6183):516–519.

2046 Loladze, I. 2002. Rising atmospheric CO₂ and human nutrition: Toward globally imbalanced
 2047 plant stoichiometry? *Trends in Ecology and Evolution*. 17(10):457–461.

2048 Loladze, I. 2014. Hidden shift of the ionome of plants exposed to elevated CO₂ depletes
 2049 minerals at the base of human nutrition. *eLife*. 2014(3):1–29.

2050 Long, S.P., Ainsworth, E.A., Rogers, A. & Ort, D.R. 2004. RISING ATMOSPHERIC
 2051 CARBON DIOXIDE: Plants FACE the Future. *Annual Review of Plant Biology*.

- 2052 55(1):591–628.
- 2053 Luo, J., Li, H., Liu, T., Polle, A., Peng, C. & Luo, Z. Bin. 2013. Nitrogen metabolism of two
 2054 contrasting poplar species during acclimation to limiting nitrogen availability. *Journal*
 2055 *of Experimental Botany*. 64(14):4207–4224.
- 2056 Madhavan, S., Treichel, I. & O’Leary, M.H. 1991. Effects of Relative Humidity on Carbon
 2057 Isotope Fractionation in Plants. *Botanica Acta*. 104(4):292–294.
- 2058 Mae, T. 1997. Physiological nitrogen efficiency in rice: Nitrogen utilization, photosynthesis,
 2059 and yield potential. *Plant and Soil*. 196:201-210
- 2060 Makino, A. 2003. Rubisco and nitrogen relationships in rice: Leaf photosynthesis and plant
 2061 growth. *Soil Science and Plant Nutrition*. 49(3):319–327.
- 2062 Manderscheid, R., Erbs, M. & Weigel, H.J. 2014. Interactive effects of free-air CO₂
 2063 enrichment and drought stress on maize growth. *European Journal of Agronomy*.
 2064 52:11–21.
- 2065 Markelz, R.J.C., Strellner, R.S. & Leakey, A.D.B. 2011. Impairment of C₄ photosynthesis by
 2066 drought is exacerbated by limiting nitrogen and ameliorated by elevated [CO₂] in
 2067 maize. *Journal of Experimental Botany*. 62(9):3235–3246.
- 2068 Marschner, H. 1995. *Mineral nutrition of higher plants*. San Diego. Academic Press.
- 2069 Marschner, H. & Marschner, P. 2012. *Marschner’s mineral nutrition of higher plants*.
 2070 London: Academic Press.
- 2071 Marschner, P. & Rengel, Z. 2012. Nutrient Availability in Soils. *Marschner’s mineral*
 2072 *nutrition of higher plants*. 315-330.
- 2073 McDonald, E.P., Erickson, J.E. & Kruger, E.L. 2002. Can decreased transpiration limit plant

2074 nitrogen acquisition in elevated CO₂? *Functional Plant Biology*. 29(9):1115–1120.

2075 Mcgrath, J.M. & Lobell, D.B. 2013. Reduction of transpiration and altered nutrient allocation
2076 contribute to nutrient decline of crops grown in elevated CO₂ concentrations. *Plant,*
2077 *Cell and Environment*. 36(3):697–705.

2078 McMurtrie, R.E., Norby, R.J., Medlyn, B.E., Dewar, R.C., Pepper, D.A., Reich, P.B. &
2079 Barton, C.V.M. 2008. Why is plant-growth response to elevated CO₂ amplified when
2080 water is limiting, but reduced when nitrogen is limiting? A growth-optimisation
2081 hypothesis. *Functional Plant Biology*. 35(6):521–534.

2082 Moore, P.H. & Botha, F.C. 2014. *Sugarcane Physiology Biochemistry & Functional Biology*.

2083 Morison, J.I.L. & Gifford, R.M. 1983. Stomatal Sensitivity to Carbon Dioxide and Humidity.
2084 *Plant Physiology*. 71(4):789–796.

2085 Mortensen, L.M. 1987. Review: CO₂ enrichment in greenhouses. Crop Responses. *Scientia*
2086 *Horticulturae*. 33(1–2):1–25.

2087 Mott, K.A. & Parkhurst, D.F. 1991. Stomatal responses to humidity in air and helox. *Plant,*
2088 *Cell & Environment*. 14(5):509–515.

2089 Mu, Q., Zhao, M., Heinsch, F.A., Liu, M., Tian, H. & Running, S.W. 2007. Evaluating water
2090 stress controls on primary production in biogeochemical and remote sensing based
2091 models. *Journal of Geophysical Research: Biogeosciences*. 112(1):1–13.

2092 Myers, S.S., Zanolatti, A., Kloog, I., Huybers, P., Leakey, A.D.B., Bloom, A.J., Carlisle, E.,
2093 Dietterich, L.H., Fitzgerald, G., Toshihiro, H., Holbrook, N. M., Nelson, R.L.,
2094 Ottman, M.J., Raboy, V., Sakai, H., Sartor, K.A., Schwartz, J., Seneweera, S., Tausz,
2095 M., Usui, Y.. 2014. Increasing CO₂ threatens human nutrition. *Nature*.
2096 510(7503):139–42.

- 2097 Ntanos, D.A. & Koutroubas, S.D. 2002. Dry matter and N accumulation and translocation for
2098 Indica and Japonica rice under Mediterranean conditions. *Field Crops Research*.
2099 74(1):93–101.
- 2100 Oliveira, E.M.M., Ruiz, H.A., Alvarez V, V.H., Ferreira, P.A., Costa, F.O. & Almeida, I.C.C.
2101 2010. Nutrient supply by mass flow and diffusion to maize plants in response to soil
2102 aggregate size and water potential. *Revista Brasileira de Ciência do Solo*. 34(2):317–
2103 328.
- 2104 Oren, R., Sperry, J.S., Katul, G.G., Pataki, D.E., Ewers, B.E., Phillips, N. & Schäfer, K.V.R.
2105 1999. Survey and synthesis of intra- and interspecific variation in stomatal sensitivity
2106 to vapour pressure deficit. *Plant, Cell and Environment*. 22(12):1515–1526.
- 2107 Oyewole, O.A., Inselsbacher, E. & Näsholm, T. 2014. Direct estimation of mass flow and
2108 diffusion of nitrogen compounds in solution and soil. *New Phytologist*. 201(3):1056–
2109 1064.
- 2110 Papakosta, D.K. & Gagianas, A.A. 1991. Nitrogen and Dry Matter Accumulation,
2111 Remobilization, and Losses for Mediterranean Wheat during Grain Filling. *Agronomy*
2112 *Journal*. 83(5):864.
- 2113 Parts, K., Tedersoo, L., Lõhmus, K., Kupper, P., Rosenthal, K., Sõber, A. & Ostonen, I.
2114 2013. Increased air humidity and understory composition shape short root traits and
2115 the colonizing ectomycorrhizal fungal community in silver birch stands. *Forest*
2116 *Ecology and Management*. 310:720–728.
- 2117 Paterson, E., Sim, A., Standing, D., Dorward, M. & McDonald, A.J.S. 2006. Root exudation
2118 from *Hordeum vulgare* in response to localized nitrate supply. *Journal of*
2119 *Experimental Botany*. 57(10):2413–2420.

- 2120 Peirson, D.R. & Elliott, J.R. 1988. Effect of Nitrite and Bicarbonate on Nitrite Utilization in
 2121 Leaf Tissue of Bush Bean (*Phaseolus vulgaris*). *Journal of Plant Physiology*.
 2122 133(4):425–429.
- 2123 Pleijel, H., Mortensen, L., Fuhrer, J., Ojanperä, K. & Danielsson, H. 1999. Grain protein
 2124 accumulation in relation to grain yield of spring wheat (*Triticum aestivum* L.) grown
 2125 in open-top chambers with different concentrations of ozone, carbon dioxide and
 2126 water availability. *Agriculture, Ecosystems and Environment*. 72(3):265–270.
- 2127 Pleijel, H., Broberg, M.C., Högy, P. & Uddling, J. 2019. Nitrogen application is required to
 2128 realize wheat yield stimulation by elevated CO₂ but will not remove the CO₂-induced
 2129 reduction in grain protein concentration. *Global Change Biology*. 25(5):1868–1876.
- 2130 Poorter, H., Van Berkel, Y., Baxter, R., Den Hertog, J., Dijkstra, P., Gifford, R.M., Griffin,
 2131 K.L., Roumet, C., Wong, S.C. 1997. The effect of elevated CO₂ on the chemical
 2132 composition and construction costs of leaves of 27 C₃ species. *Plant, Cell and*
 2133 *Environment*. 20(4):472–482.
- 2134 Porteous, F., Hill, J., Ball, A.S., Pinter, P.J., Kimball, B.A., Wall, G.W., Adamsen, F.J.,
 2135 Hunsaker, D.J., LaMorte, R.L., Leavitt, S.W., Thompson, T.L., Matthias, A.D.,
 2136 Brooks, T.J., Morris, C.F. 2009. Effect of Free Air Carbon dioxide Enrichment
 2137 (FACE) on the chemical composition and nutritive value of wheat grain and straw.
 2138 *Animal Feed Science and Technology*. 149(3–4):322–332.
- 2139 del Pozo, A., Perez, P., Gutierrez, D., Alonso, A., Morcuende, R. & Martinez-Carrasco, R.
 2140 2007. Gas exchange acclimation to elevated CO₂ in upper-sunlit and lower-shaded
 2141 canopy leaves in relation to nitrogen acquisition and partitioning in wheat grown in
 2142 field chambers. *Environmental and Experimental Botany*. 59:371–380.
- 2143 Pritchard, S. & Rogers, H. 2000. Spatial and temporal deployment of crop. *New Phytologist*.

2144 147:55–71.

2145 R Core Team (2019). R: A language and environment for statistical computing. R Foundation
 2146 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

2147 Ranathunge, K., Steudle, E. & Lafitte, R. 2003. Control of water uptake by rice (*Oryza sativa*
 2148 L.): role of the outer part of the root. *Planta*. 217:193–205.

2149 Rathnam, C.K.M. 1978. Malate and Dihydroxyacetone Phosphate-dependent Nitrate
 2150 Reduction in Spinach Leaf Protoplasts. *Plant Physiology*. 62(2):220–223.

2151 Reeves, D.W., Rogers, H.H., Prior, S.A., Wood, C.W. & Runion, G.B. 1994. Elevated
 2152 atmospheric carbon dioxide effects on sorghum and soybean nutrient status. *Journal*
 2153 *of Plant Nutrition*. 17(11):1939–1954.

2154 Reich, P.B., Walters, M.B. & Ellsworth, D.S. 1997. From tropics to tundra: global
 2155 convergence in plant functioning. *Proceedings of the National Academy of Sciences*.
 2156 94:13730–13734.

2157 Robinson, S.P. & Giersch, C. 1987. Inorganic Phosphate Concentration in the Stroma of
 2158 Isolated Chloroplasts and Its Influence on Photosynthesis. *Australian Journal of Plant*
 2159 *Physiology*. 14(4):451–462

2160 Rogers, H.H. & Runion, G.B. 1994. Plant responses to atmospheric CO₂ enrichment with
 2161 emphasis on roots and the rhizosphere. *Environmental Pollution*. 83:155–189.

2162 Rose, L., Buitenwerf, R., Cramer, M., February, E.C. & Higgins, S.I. 2018. Effects of
 2163 nutrient supply on carbon and water economies of C₄ grasses. *Functional Plant*
 2164 *Biology*. 45(9):935–944.

2165 Rosenvald, K., Tullus, A., Ostonen, I., Uri, V., Kupper, P., Aosaar, J., Varik, M., Söber, J.,
 2166 Niglas, A., Hansen, R., Rohula, G., Kukk, M., Sober, A., Lohmus, K. 2014. The effect

2167 of elevated air humidity on young silver birch and hybrid aspen biomass allocation
 2168 and accumulation - acclimation mechanisms and capacity. *Forest Ecology and*
 2169 *Management*. 330:252–260.

2170 Rufty, T.W., Thomas, J.F., Remmler, J.L., Campbell, W.H. & Volk, R.J. 1986. Intercellular
 2171 Localization of Nitrate Reductase in Roots. *Plant Physiology*. 82(3):675–680.

2172 Russell, R.S. & Barber, S.A. 1960. The Relationship Between Salt Uptake and the
 2173 Absorption of Water by Intact Plants. *Annual Review of Plant Physiology*. 11:127–
 2174 140.

2175 Sage, R.F. 1994. Acclimation of photosynthesis to increasing atmospheric CO₂: The gas
 2176 exchange perspective. *Photosynthesis Research*. 39(3):351–368.

2177 Sanchez-Diaz, M., Garcia, J.L., Antolin, M.C. & Araus, J.L. 2002. Effects of soil drought and
 2178 atmospheric humidity on yield, gas exchange, and stable carbon isotope composition
 2179 of barley. *Photosynthetica*. 40(3):415–421.

2180 Schmidt, I.K., Tietema, A., Williams, D., Gundersen, P., Beier, C., Emmett, B.A. & Estiarte,
 2181 M. 2004. Soil Solution Chemistry and Element Fluxes in Three European Heathlands
 2182 and Their Responses to Warming and Drought. *Ecosystems*. 7(6):638–649.

2183 Searles, P.S. & Bloom, A.J. 2003. Nitrate photo-assimilation in tomato leaves under short-
 2184 term exposure to elevated carbon dioxide and low oxygen. 1247–1255.

2185 Sellin, A., Tullus, A., Niglas, A., Õunapuu, E., Karusion, A. & Lõhmus, K. 2013. Humidity-
 2186 driven changes in growth rate, photosynthetic capacity, hydraulic properties and other
 2187 functional traits in silver birch (*Betula pendula*). *Ecological Research*. 28(3):523–535.

2188 Sellin, A., Rosenvald, K., Õunapuu-Pikas, E., Tullus, A., Ostonen, I. & Lõhmus, K. 2015.
 2189 Elevated air humidity affects hydraulic traits and tree size but not biomass allocation

2190 in young silver birches (*Betula pendula*). *Frontiers in Plant Science*. 6:1–11.

2191 Sharkey, T.D. 1984. Transpiration-induced changes in the photosynthetic capacity of leaves.
2192 *Planta*. 160(2):143–150. DOI: 10.1007/BF00392862.

2193 Simpson, R.J., Lambers, H. & Dalling, M.J. 1983. Nitrogen Redistribution during Grain
2194 Growth in Wheat (*Triticum aestivum* L.) . *Plant Physiology*. 71(1):7–14.

2195 Sinclair, T.R., Zwieniecki, M.A. & Holbrook, N.M. 2008. Low leaf hydraulic conductance
2196 associated with drought tolerance in soybean. *Physiologia Plantarum*. 132(4):446–
2197 451.

2198 Singh, S. 2016. Guttation: Mechanism, Momentum and Modulation. *Botanical Review*.
2199 82(2):149–182.

2200 Slewinski, T.L., Meeley, R. & Braun, D.M. 2009. Sucrose transporter1 functions in phloem
2201 loading in maize leaves. *Journal of Experimental Botany*. 60(3):881–892.

2202 Smith, M.R. & Myers, S.S. 2019. Global Health Implications of Nutrient Changes in Rice
2203 Under High Atmospheric Carbon Dioxide. *GeoHealth*. 3(7):190–200 .

2204 Smith, P., Davis, S.J., Creutzig, F., Fuss, S., Minx, J., Gabrielle, B., Kato, E., Jackson, R.B.,
2205 et al. 2016. Biophysical and economic limits to negative CO₂ emissions. *Nature*
2206 *Climate Change*. 6(1):42–50.

2207 Stimler, K., Berry, J.A. & Yakir, D. 2012. Effects of carbonyl sulfide and carbonic anhydrase
2208 on stomatal conductance. *Plant Physiology*. 158(1):524–530.

2209 Stitt, M. & Krapp, A. 1999. The interaction between elevated carbon dioxide and nitrogen
2210 nutrition: the physiological and molecular background. *Plant, Cell and Environment*.
2211 22:553–621.

- 2212 Taiz, L. & Zeiger, E. 2010. *Plant Physiology (Fifth Edition)*. V. 1.
- 2213 Tanner, W. & Beevers, H. 2001. Transpiration, a prerequisite for long-distance transport of
 2214 minerals in plants? *Proceedings of the National Academy of Sciences of the United*
 2215 *States of America*. 98(16):9443–9447.
- 2216 Tanner, W. & Beevers, H. 1990. Does transpiration have an essential function in long-
 2217 distance ion transport in plants? *Plant, Cell & Environment*. 13(8):745–750.
- 2218 Taub, D.R. & Wang, X. 2008. Why are nitrogen concentrations in plant tissues lower under
 2219 elevated CO₂? A critical examination of the hypotheses. *Journal of Integrative Plant*
 2220 *Biology*. 50(11):1365–1374.
- 2221 Taub, D.R., Miller, B. & Allen, H. 2008. Effects of elevated CO₂ on the protein concentration
 2222 of food crops: a meta-analysis. *Global Change Biology*. 14(3):565–575. DOI:
 2223 10.1111/j.1365-2486.2007.01511.x.
- 2224 Tullus, A., Kupper, P., Sellin, A., Parts, L., Söber, J., Tullus, T., Löhmus, K., Söber, A.,
 2225 Tullus, H. 2012. Climate change at Northern latitudes: Rising atmospheric humidity
 2226 decreases transpiration, N-uptake and growth rate of hybrid aspen. *PLoS ONE*. 7(8):
 2227 e42648
- 2228 Turner, N.C., Schulze, E. & Gollan, T. 1985. The Responses of Stomata and Leaf Gas
 2229 Exchange to Vapour Pressure Deficits and Soil Water Content . II. In the Mesophytic
 2230 Herbaceous Species *Helianthus annuus*. *Oecologia*. 65(3):348–355.
- 2231 Uddling, J., Broberg, M.C., Feng, Z. & Pleijel, H. 2018. Crop quality under rising
 2232 atmospheric CO₂. *Current Opinion in Plant Biology*. 45:262–267.
- 2233 Vandamme, E., Renkens, M., Pypers, P., Smolders, E., Vanlauwe, B. & Merckx, R. 2013.
 2234 Root hairs explain P uptake efficiency of soybean genotypes grown in a P-deficient

- 2235 Ferralsol. *Plant and Soil*. 369(1–2):269–282.
- 2236 Van Vuuren, M.M.I., Robinson, D., Fitter, A.H., Chasalow, S.D., Williamson, L. & Raven,
2237 J.A. 1997. Effects of elevated atmospheric CO₂ and soil water availability on root
2238 biomass, root length, and N, P and K uptake by wheat. *New Phytologist*. 135(3):455–
2239 465.
- 2240 Wallace, M.B. & Knausenberger, W.I. 1997. Inorganic fertilizer use in africa :
2241 *Environmental and economic dimensions*. 10–13
- 2242 Wand, S.J.E., Midgley, G.F., Jones, M.H. & Curtis, P.S. 1999. Responses of wild C₄ and C₄
2243 grass (Poaceae) species to elevated atmospheric CO₂ concentrations: a meta-analytic
2244 test of current theories and perceptions. *Global Change Biology*. 5:723–741.
- 2245 Webber, A.N., Nie, G.Y. & Long, S.P. 1994. Acclimation of photosynthetic proteins to rising
2246 atmospheric CO₂. *Photosynthesis Research*. 39(3):413–425.
- 2247 Weltzin, J.F., Loik, M.E., Schwinning, S., Williams, D.G., Fay, P.A., Haddad, B.M., Harte,
2248 J., Huxman, T.E., Knapp, A.K., Lin, G., Pockman, W.T., Shaw, R.M., Small, E.E.,
2249 Smith, M.D., Smith, S.D., Tissue, D.T., Zak, J.C. 2003. Assessing the Response of
2250 Terrestrial Ecosystems to Potential Changes in Precipitation. *BioScience*. 53(10):941.
- 2251 Wen, Z., Li, H., Shen, J. & Rengel, Z. 2017. Maize responds to low shoot P concentration by
2252 altering root morphology rather than increasing root exudation. *Plant and Soil*. 416(1–
2253 2):377–389.
- 2254 H. Wickham. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York, 2016.
- 2255 Winter, K., Holtum, J.A.M., Edwards, G.E. & O’leary, M.H. 1982. Effect of low relative
2256 humidity on $\delta^{13}\text{C}$ value in two C₃ grasses and in *Panicum milioides*, a C₃-C₄
2257 intermediate species. *Journal of Experimental Botany*. 33(1):88–91.


- 2258 Wong, S.C. 1979. Elevated atmospheric partial pressure of CO₂ and plant growth - I.
2259 Interactions of nitrogen nutrition and photosynthetic capacity in C₃ and C₄ plants.
2260 *Oecologia*. 44(1):68–74.
- 2261 Wong, S.C. 1990. Elevated atmospheric partial pressure of CO₂ and plant growth - II. Non-
2262 structural carbohydrate content in cotton plants and its effect on growth parameters.
2263 *Photosynthesis Research*. 23(2):171–180.
- 2264 Woodward, R.G. & Begg, J.E. 1976. The effect of atmospheric humidity on the yield of
2265 soybean. *Australian Journal of Agricultural Research*. 27(4):501–508.
- 2266 Wright, G.C., Rao, R.C.N. & Farquhar, G.D. 1994. Water-Use Efficiency and Carbon Isotope
2267 Discrimination in Peanut under Water Deficit Conditions. *Crop Science*. 34:92–97.
- 2268 Xu, G., Fan, X. & Miller, A.J. 2012. Plant Nitrogen Assimilation and Use Efficiency. *Annual*
2269 *Review of Plant Biology*. 63(1):153–182.
- 2270 Xue, Q., Weiss, A., Arkebauer, T.J. & Baenziger, P.S. 2004. Influence of soil water status
2271 and atmospheric vapor pressure deficit on leaf gas exchange in field-grown winter
2272 wheat. *Environmental and Experimental Botany*. 51(2):167–179.
- 2273 Yang, Z., Sinclair, T.R., Zhu, M., Messina, C.D., Cooper, M. & Hammer, G.L. 2012.
2274 Temperature effect on transpiration response of maize plants to vapour pressure
2275 deficit. *Environmental and Experimental Botany*. 78:157–162.
- 2276 Zhao, F.J. & McGrath, J.M. 2002. Assessing bioavailability of zinc, cadmium and copper in
2277 soils to the indicator plant *Lepidium heterophyllum*. In *COST Action 837. 4th WG2*
2278 *Workshop, Bordeaux'2002 – Risk assessment and sustainable land management using*
2279 *plants in trace element-contaminated soils*. Bordeaux.
- 2280 Zhu, C., Kobayashi, K., Loladze, I., Zhu, J., Jiang, Q., Xu, X., Liu, G., Seneweera, S., Ebi,

2281 K.L., Drewnowski, A., Fukagawa, N.K., Ziska, L.H. 2018. Carbon dioxide (CO₂)
2282 levels this century will alter the protein, micronutrients and vitamin content of rice
2283 grains with potential health consequences for the poorest rice-dependent countries.
2284 *Science Advances*. 4:1–9.

2285 Ziska, L.H. & Bunce, J.A. 1997. Influence of increasing carbon dioxide concentration on the
2286 photosynthetic and growth stimulation of selected C₄ crops and weeds. *Photosynthesis*
2287 *Research*. 54(3):199–208.

2288

2289



2291

2292 **Appendix 1** Variation in transpiration-driven mass flow delivery of Ca, Cu and Mn in maize

2293 and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels,

2294 400 and 800 ppm. The boxes and horizontal lines represent the first and the third quartiles and


2295 the medians, respectively. The whiskers represent 1.5 × the interquartile range and outliers

2296 above/below are shown as points. The diamonds represent the mean values. The F- and p-


2297 values of the main effects and interaction (when significant) from the analysis of variance done

2298 on the linear models fitted are displayed. Lower case letters indicate the significant differences


2299 between treatments as determined by Tukey's post-hoc comparisons.




Appendix 2 Variation in transpiration-driven mass flow delivery of Fe and Zn in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.




Appendix 3 Variation in transpiration-driven mass flow contribution of Ca, Cu and Mn to tissue content in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The dashed line represents 100% contribution to tissue content. Statistics as in Appendix 1.




Appendix 4 Variation in transpiration-driven mass flow contribution of Fe and Zn to tissue content in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The dashed line represents 100% contribution to tissue content. Statistics as in Appendix 1.




Appendix 5 Variation in total tissue concentrations of Ca, Cu and Mn in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO_2 levels, 400 and 800 ppm. Statistics as in Appendix 1.




Appendix 6 Variation in total tissue concentrations of Fe and Zn in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.




Appendix 7 Variation in total tissue contents of Ca, Cu and Mn in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.




Appendix 8 Variation in total tissue contents of Fe and Zn in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.




Appendix 9 The relationship between transpiration-driven mass flow delivery of Ca, Cu and Mn, and total tissues contents of Ca, Cu and Mn, in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. The intercept, slope, R² and p-values of the lines are displayed, as well as the p-value of the contrast between the 400 and 800 ppm line.




Appendix 10 The relationship between transpiration-driven mass flow delivery of Fe and Zn, and total tissues contents of Fe and Zn, in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 9.



Appendix 11 Variation in the C:Ca, C:Cu and C:Mn ratios in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.



Appendix 12 Variation in the C:Fe and C:Zn ratios in maize and wheat plants, grown in five soils with differing nutrient availabilities at two CO₂ levels, 400 and 800 ppm. Statistics as in Appendix 1.



2363

2364

2365

2366

2367

2368


2369

2370


2371

2372

Appendix 13 Variation in transpiration-driven mass flow delivery of Ca, Cu and Mn in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. The boxes and horizontal lines represent the first and the third quartiles and the medians, respectively. The whiskers represent $1.5 \times$ the interquartile range and outliers above/below are shown as points. The diamonds represent the mean values. The F- and p-values of the main effects and interaction (when significant/present) from the analysis of variance done on the linear models fitted are displayed. Lower case letters indicate the significant differences between treatments as determined by Tukey's post-hoc comparisons. Corresponding air relative humidity (in %) at 25°C shown on the x-axis.




Appendix 14 Variation in transpiration-driven mass flow delivery of Fe and Zn in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Appendix 1. Corresponding air relative humidity (in %) at 25°C shown on the x-axis.




2378

2379 **Appendix 15** Variation in transpiration-driven mass flow contribution of Ca, Cu and Mn to
 2380 tissue content in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa,
 2381 0.773 kPa and 0.350 kPa. The dashed line represents 100% contribution to tissue content.
 2382 Statistics as in Appendix 1. Corresponding air relative humidity (in %) at 25°C shown on the
 2383 x-axis.


2384



Appendix 16 Variation in transpiration-driven mass flow contribution of Fe and Zn to tissue content in maize and wheat plants, grown in Soil, at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. The dashed line represents 100% contribution to tissue content. Statistics as in Appendix 1. Corresponding air relative humidity (in %) at 25°C shown on the x-axis.




Appendix 17 Variation in total tissue concentrations of Ca, Cu and Mn in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Appendix 1.




2396

2397 **Appendix 18** Variation in total tissue concentrations of Fe and Zn in maize and wheat plants,
 2398 grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics
 2399 as in Appendix 1.


2400



Appendix 19 Variation in total tissue contents of N, P and K in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Appendix 1.



Appendix 20 Variation in total tissue contents of Ca, Cu and Mn in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Appendix 1.



Appendix 21 Variation in total tissue contents of Fe and Zn in maize and wheat plants, grown in Soil and in Sand at three VPD levels, 1.613 kPa, 0.773 kPa and 0.350 kPa. Statistics as in Appendix 1.